

A TOPOLOGICAL PROPERTY OF ASYMPTOTICALLY CONICAL SELF-SHRINKERS OF SMALL ENTROPY

JACOB BERNSTEIN AND LU WANG

ABSTRACT. For any asymptotically conical self-shrinker with entropy less than or equal to that of a cylinder we show that the link of the asymptotic cone must separate the unit sphere into exactly two connected components, both diffeomorphic to the self-shrinker. Combining this with recent work of Brendle, we conclude that the round sphere uniquely minimizes the entropy among all non-flat two-dimensional self-shrinkers. This confirms a conjecture of Colding-Ilmanen-Minicozzi-White in dimension two.

1. INTRODUCTION

A hypersurface $\Sigma \subset \mathbb{R}^{n+1}$ is said to be a *self-shrinker* if it satisfies

$$(1.1) \quad \mathbf{H}_\Sigma + \frac{\mathbf{x}^\perp}{2} = \mathbf{0}.$$

Here $\mathbf{H}_\Sigma = -H_\Sigma \mathbf{n}_\Sigma = \Delta_\Sigma \mathbf{x}$ is the mean curvature vector of Σ and $\mathbf{x}^\perp$ is the normal component of the position vector. Self-shrinkers arise naturally in the mean curvature flow as the time slices of solutions that move self-similarly by scaling. Specifically, if Σ is a self-shrinker, then

$$(1.2) \quad \{\Sigma_t\}_{t \in (-\infty, 0)} = \{\sqrt{-t}\Sigma\}_{t \in (-\infty, 0)}$$

is a smooth mean curvature flow. Such self-similar flows are important as they model singularities of the mean curvature flow. From a variational point of view, self-shrinkers arise as critical points of the *Gaussian (hyper)-surface area*

$$(1.3) \quad F[\Sigma] = \int_\Sigma \Phi(\mathbf{x}) d\mathcal{H}^n = (4\pi)^{-\frac{n}{2}} \int_\Sigma e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n.$$

Here $\mathcal{H}^n$ is n -dimensional Hausdorff measure and Φ is the Gaussian normalized so that $F[\mathbb{R}^n \times \{0\}] = 1$. Following Colding-Minicozzi [10], the *entropy* of a hypersurface is defined by

$$(1.4) \quad \lambda[\Sigma] = \sup_{(\mathbf{y}, \rho) \in \mathbb{R}^{n+1} \times \mathbb{R}^+} F[\rho\Sigma + \mathbf{y}].$$

The Gaussian surface area and the entropy of self-shrinkers agree and so $\lambda[\mathbb{R}^n \times \{0\}] = 1$.

A hypersurface, Σ , is *asymptotically conical*, if it is smoothly asymptotic to a regular cone; i.e., $\lim_{\rho \rightarrow 0} \rho\Sigma = \mathcal{C}(\Sigma)$ in $C_{loc}^\infty(\mathbb{R}^{n+1} \setminus \{0\})$ for $\mathcal{C}(\Sigma)$ a regular cone. Our main result is a topological restriction on asymptotically conical self-shrinkers with small entropy:

Theorem 1.1. *Let Σ be an asymptotically conical self-shrinker in $\mathbb{R}^{n+1}$ for $n \geq 2$. If $\lambda[\Sigma] \leq \lambda[\mathbb{S}_*^{n-1}]$, then $\mathcal{L}(\Sigma)$, the link of the asymptotic cone $\mathcal{C}(\Sigma)$, separates $\mathbb{S}^n$ into two connected components both diffeomorphic to Σ . As a consequence, $\mathcal{L}(\Sigma)$ is connected.*

The first author was partially supported by the NSF Grant DMS-1307953. The second author was partially supported by the Chapman Fellowship of the Imperial College London and by the AMS-Simons Travel Grant 2012-2014 and the NSF Grant DMS-1406240.

Here $\mathbb{S}^n \subset \mathbb{R}^{n+1}$ is the unit sphere centered at the origin and, for $0 \leq k \leq n$,

$$(1.5) \quad \mathbb{S}_*^{n-k} \times \mathbb{R}^k = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^{n-k+1} \times \mathbb{R}^k = \mathbb{R}^{n+1} : |\mathbf{x}|^2 = 2(n-k)\}$$

are the maximally symmetric self-shrinking cylinders with k -dimensional spine¹. As the $\mathbb{S}_*^{n-k} \times \mathbb{R}^k$ are self-shrinkers, their Gaussian surface area and entropy agree. That is,

$$(1.6) \quad \lambda_n = \lambda[\mathbb{S}^n] = F[\mathbb{S}_*^n] = F[\mathbb{S}_*^n \times \mathbb{R}^l].$$

By a computation of Stone [24],

$$(1.7) \quad 2 > \lambda_1 > \frac{3}{2} > \lambda_2 > \dots > \lambda_n > \dots \rightarrow \sqrt{2}.$$

When $n = 2$, Theorem 1.1 implies that Σ is diffeomorphic to an open disk. This allows us to completely classify self-shrinkers in $\mathbb{R}^3$ of small entropy. Indeed, combining the work of Colding-Ilmanen-Minicozzi-White [9] (see also [2]) and Brendle [4] with Theorem 1.1, we conclude:

Corollary 1.2. *If $\Sigma \subset \mathbb{R}^3$ is a self-shrinker with $\lambda[\Sigma] \leq \lambda_1$, then Σ is either $\mathbb{S}_*^2$, or some rotation of $\mathbb{R}^2 \times \{0\}$ or $\mathbb{S}_*^1 \times \mathbb{R}$. In fact, there is a $\delta_0 > 0$ so that these are the only self-shrinkers with entropy less than $\lambda_1 + \delta_0$.*

This answers affirmatively the $n = 2$ case of [9, Conjecture 0.10] which asks whether $\mathbb{S}_*^n$ minimizes the entropy amongst all non-flat self-shrinkers. In addition, this gives, for mean curvature flows in $\mathbb{R}^3$, that λ_2 is the best constant in White's formulation [26] of Brakke's regularity theorem [3]. Another simple consequence of Corollary 1.2 is a sharp lower bound for the entropy of all closed surfaces in $\mathbb{R}^3$ (see [2, Theorem 1.1] for a different approach which works for all $2 \leq n \leq 6$) and a new entropy lower bound for all closed surfaces in $\mathbb{R}^3$ of positive genus:

Corollary 1.3. *If Σ is a closed surface in $\mathbb{R}^3$, then $\lambda[\Sigma] \geq \lambda_2$ with equality if and only if Σ is, up to translations and scalings, $\mathbb{S}_*^2$. Furthermore, if Σ has positive genus, then $\lambda[\Sigma] > \lambda_1$.*

Our proof of Theorem 1.1 parallels that of Colding-Ilmanen-Minicozzi-White, who, in [9], showed that $\mathbb{S}_*^n$ uniquely minimizes the entropy within the class of closed self-shrinkers. Indeed, our arguments may be thought of as a natural analog, in the asymptotically conical setting, of their arguments. However, we wish to emphasize one crucial difference between our proofs. Namely, in [9] the authors exploit the fact that a closed hypersurface must form a singularity in finite time – indeed, they point out this is the main obstruction to extending their result to the non-compact setting. We, however, make use of the fact that flows of asymptotically conical surfaces of small entropy must exist without singularities for long-time to show that the flows eventually become star-shaped with respect to the origin.

More precisely, inspired by [9], we first study properties of a quantity along a smooth mean curvature flow that we call the *shrinker mean curvature* – see (3.1). In particular, we use a parabolic maximum principle on non-compact manifolds as in [12] to show that if the initial surface of a smooth mean curvature flow is well-behaved at infinity and has positive shrinker mean curvature, then the shrinker mean curvature remains positive along the flow – see Proposition 3.2. We also use a standard parabolic maximum principle to conclude

¹In $\mathbb{R}^{n+1}$ a round sphere is a self-shrinker if and only if it is centered at the origin and its radius is $\sqrt{2n}$. In our notation, the radii of $\mathbb{S}_*^n$ and $\mathbb{S}^n$ are, respectively, $\sqrt{2n}$ and 1. Thus, $\mathbb{S}_*^n$ is a self-shrinker while $\mathbb{S}^n$ is not.

that for such a flow, the shrinker mean curvature controls the second fundamental form – see Proposition 3.3.

A variant on the shrinker mean curvature is also considered in [9] where the above properties are shown to hold for smooth mean curvature flows of *closed* hypersurfaces. However, in [9] the authors find it convenient to think of self-shrinkers as static points and so instead work with a certain rescaled mean curvature flow. As such, their quantity has a different, but related, form. While this may appear to be merely a technical distinction it actually holds the key to our proof. Indeed, the quantity we consider makes sense for smooth mean curvature flows which start close to a self-shrinker but that persist up to (and beyond) the singular time of the self-shrinker. By its construction, this cannot be true of the quantity considered in [9].

We next observe that if Σ is any non-flat asymptotically conical self-shrinker, then there are asymptotically conical perturbations $\Gamma^\pm$ of Σ on each side of Σ which have positive shrinker mean curvature (relative to the correct orientation), strictly smaller entropy and which have the same asymptotic cone as that of Σ – see Proposition 4.2. As in [9], these two hypersurfaces are found by considering the normal exponential graphs of small multiples of the lowest eigenfunction of the (self-shrinker) stability operator. Again the non-compactness introduces certain technical difficulties.

In our third step, we consider the smooth mean curvature flows with initial surfaces $\Gamma^\pm$. Directly using arguments in [9], we prove that for non-compact flows of positive shrinker mean curvature if a singularity develops, then the Gaussian density of the flows at the singular point must be at least λ_{n-1} . Hence, if the entropy of Σ is at most λ_{n-1} , then the flows starting from $\Gamma^\pm$ never develop singularities. We further show that each time slice of these flows is smoothly asymptotic to the same cone as that of Σ .

Finally, we observe that any long-time solution of the mean curvature flow with positive shrinker mean curvature and which is asymptotically conical must have a time-slice which is star-shaped – see Theorem 4.6 and Proposition 5.1. Theorem 1.1 is then an immediate consequence. The proofs of Corollaries 1.2 and 1.3 are straightforward.

2. NOTATION

Let $\mathbb{R}^{n+1}$ denote the standard $(n+1)$ -dimensional Euclidean space. We denote by

$$(2.1) \quad B_R(\mathbf{x}_0) = \{\mathbf{x} \in \mathbb{R}^{n+1} : |\mathbf{x} - \mathbf{x}_0| < R\}$$

the open Euclidean ball of radius R centered at $\mathbf{x}_0$. When the center is the origin $\mathbf{0}$, we will omit it. A *hypersurface* in $\mathbb{R}^{n+1}$ is a proper codimension one submanifold of $\mathbb{R}^{n+1}$. At times it will be convenient to distinguish between a point $p \in \Sigma$ and its *position vector* $\mathbf{x}(p) \in \mathbb{R}^{n+1}$. Given a set $\Omega \subset \mathbb{R}^{n+1}$ we define the ρ -tubular neighborhood of Ω to be the open set

$$(2.2) \quad \mathcal{T}_\rho(\Omega) = \bigcup_{p \in \Omega} B_\rho(p) \subset \mathbb{R}^{n+1}.$$

Any hypersurface is two-sided and hence orientable – see for instance [20]. For this reason, we will always orient a hypersurface, Σ , by a choice of unit normal

$$(2.3) \quad \mathbf{n}_\Sigma : \Sigma \rightarrow \mathbb{S}^n \subset \mathbb{R}^{n+1}.$$

For such a Σ , we let g_Σ denote the induced Riemannian metric, A_Σ be the second fundamental form (with respect to the choice of unit normal $\mathbf{n}_\Sigma$) and H_Σ the associated scalar mean curvature which we take to be the trace of A_Σ . Observe that the mean curvature vector $\mathbf{H}_\Sigma = -H_\Sigma \mathbf{n}_\Sigma$ is well-defined independent of choices of unit normal as is the norm

$|A_\Sigma|$ of the second fundamental form. For any hypersurface, Σ , and $\rho \in \mathbb{R}^+$ we have the natural smooth map

$$(2.4) \quad \Psi_\rho : \Sigma \times (-\rho, \rho) \rightarrow \mathcal{T}_\rho(\Sigma)$$

given by $\Psi_\rho(p, s) = \mathbf{x}(p) + s\mathbf{n}_\Sigma(p)$. We say $\mathcal{T}_\rho(\Sigma)$ is a *regular tubular neighborhood* if this map is a diffeomorphism.

We consider space-time, $\mathbb{R}^{n+1} \times \mathbb{R}$, to be the set of space-time points $X = (\mathbf{x}, t) \in \mathbb{R}^{n+1} \times \mathbb{R}$. And let $O = (0, 0)$ be the space-time origin. We will often focus on the subset of space-time consisting of space-time points with negative time which we denote by $\mathbb{R}^{n+1} \times \mathbb{R}^-$. Given $R, \tau > 0$, let

$$(2.5) \quad C_{R,\tau}(X_0) = \{(\mathbf{x}, t) \in \mathbb{R}^{n+1} \times \mathbb{R} : \mathbf{x} \in B_R(\mathbf{x}_0), |t - t_0| < \tau\}$$

be the parabolic cylinder of radius R and height τ centered at $X_0 = (\mathbf{x}_0, t_0)$. It will also be convenient to consider the backward parabolic cylinder

$$(2.6) \quad C_{R,\tau}^-(X_0) = C_{R,\tau}(X_0) \cap \mathbb{R}^{n+1} \times \{t < t_0\}.$$

The *parabolic boundary* of a parabolic cylinder is defined to be

$$(2.7) \quad \partial_P C_{R,\tau}(X_0) = \partial C_{R,\tau}(X_0) \setminus (B_R(\mathbf{x}_0) \times \{t = t_0 + \tau\}),$$

where ∂ is the topological boundary. Likewise, $\partial_P C_{R,\tau}^-(X_0) = \partial C_{R,\tau}^-(X_0) \cap \partial_P C_{R,\tau}(X_0)$.

A *smooth mean curvature flow* is a collection of hypersurfaces $\{\Sigma_t\}_{t \in I}$, where I is some interval in $\mathbb{R}$ and so that there is a smooth map $F : M \times I \rightarrow \mathbb{R}^{n+1}$ so that for each $t \in I$, $F(\cdot, t) : M \rightarrow \Sigma_t \subset \mathbb{R}^{n+1}$ is a parameterization of Σ_t and so that

$$(2.8) \quad \left(\frac{\partial}{\partial t} F(p, t) \right)^\perp = \mathbf{H}_{\Sigma_t}(F(p, t)).$$

We will always take the hypersurfaces Σ_t in a smooth mean curvature flow to be oriented so that the unit normal is smooth in t . It is often convenient to consider the *space-time track* of a smooth mean curvature flow

$$(2.9) \quad \mathcal{S} = \{(\mathbf{x}(p), t) \in \mathbb{R}^{n+1} \times \mathbb{R} : p \in \Sigma_t\},$$

which is a smooth submanifold of space-time (with boundary if I contains either of its endpoints) that is transverse to each constant time hyperplane. We will not distinguish between a smooth mean curvature flow $\{\Sigma_t\}_{t \in I}$ and its space-time track and so denote both by $\mathcal{S}$. Along the space-time track of a smooth mean curvature flow $\mathcal{S}$, let $\frac{d}{dt}$ be the smooth vector field given by

$$(2.10) \quad \frac{d}{dt} = \frac{\partial}{\partial t} + \mathbf{H}_{\Sigma_t}.$$

It is not hard to see that this vector field is tangent to $\mathcal{S}$ and the position vector satisfies

$$(2.11) \quad \frac{d}{dt} \mathbf{x}(p, t) = \mathbf{H}_{\Sigma_t}(p).$$

It is straightforward (and standard) to compute the evolution of various geometric quantities with respect to this vector field – see for instance [11, Appendix B].

For a set $\Omega \subset \mathbb{R}^{n+1}$, an $\mathbf{x} \in \mathbb{R}^{n+1}$ and a $\rho \in \mathbb{R}^+$, let

- (1) $\Omega + \mathbf{x} = \{\mathbf{y} \in \mathbb{R}^{n+1} : \mathbf{y} - \mathbf{x} \in \Omega\}$, the translation of Ω by $\mathbf{x}$; and
- (2) $\rho \Omega = \{\mathbf{y} \in \mathbb{R}^{n+1} : \rho^{-1} \mathbf{y} \in \Omega\}$, the scaling of Ω by ρ .

Similarly, for a set $\Omega \subset \mathbb{R}^{n+1} \times \mathbb{R}$ in space-time, an $X \in \mathbb{R}^{n+1} \times \mathbb{R}$ and a $\rho \in \mathbb{R}^+$,

- (1) $\Omega + X = \{Y \in \mathbb{R}^{n+1} \times \mathbb{R} : Y - X \in \Omega\}$, the space-time translation of Ω by X ;
 and
 (2) $\rho\Omega = \{(y, t) \in \mathbb{R}^{n+1} : (\rho^{-1}y, \rho^{-2}t) \in \Omega\}$, the parabolic scaling of Ω by ρ .

Clearly, if $\mathcal{S}$ is the space-time track of a smooth mean curvature flow, then so is $\rho\mathcal{S} + X$ for any $\rho \in \mathbb{R}^+$ and $X \in \mathbb{R}^{n+1} \times \mathbb{R}$.

3. SHRINKER MEAN-CONVEXITY

Let $\mathcal{S} = \{\Sigma_t\}_{t \in [-1, T]}$ be a smooth mean curvature flow. Along the flow $\mathcal{S}$, we define the *shrinker mean curvature relative to the space-time point* $X_0 = (\mathbf{x}_0, t_0)$ to be

$$(3.1) \quad S_{\Sigma_t}^{X_0}(p) = 2(t_0 - t)H_{\Sigma_t}(p) - (\mathbf{x}(p) - \mathbf{x}_0) \cdot \mathbf{n}_{\Sigma_t}(p).$$

We emphasize that, due to the t dependence, this quantity is defined for $\mathcal{S}$. Given a time $t \in \mathbb{R}$ and hypersurface Σ , the shrinker mean curvature of Σ relative to the space-time point X_0 and time t is defined to be

$$(3.2) \quad S_{\Sigma}^{X_0, t}(p) = 2(t_0 - t)H_{\Sigma}(p) - (\mathbf{x}(p) - \mathbf{x}_0) \cdot \mathbf{n}_{\Sigma}(p).$$

We recall the following evolution equation for the shrinker mean curvature due to Smoczyk [23, Proposition 4].

Lemma 3.1. *Along a mean curvature flow $\{\Sigma_t\}_{t \in I}$, the shrinker mean curvature relative to X_0 satisfies*

$$(3.3) \quad \frac{d}{dt} S_{\Sigma_t}^{X_0} = \Delta_{\Sigma_t} S_{\Sigma_t}^{X_0} + |A_{\Sigma_t}|^2 S_{\Sigma_t}^{X_0}.$$

Observe that $\mathcal{S}$ is self-similar with respect to parabolic rescalings about X_0 if and only if $S_{\Sigma_t}^{X_0} \equiv 0$. More generally, if one parabolically dilates $\mathcal{S}$ about X_0 , then the vector field of the normal variation of this family at $\mathcal{S}$ is the shrinker mean curvature vector relative to X_0 . Hence, as parabolic dilations are symmetries of the mean curvature flow, (3.3) may be viewed as the linearization of the mean curvature flow. In a similar fashion, the quantities $\mathbf{e}_i \cdot \mathbf{n}_{\Sigma_t}$ used by Ecker-Huisken [12] are generated by spatial translations and so also satisfy (3.3). Likewise, the mean curvature H_{Σ_t} which also satisfies (3.3) arises in this manner from temporal translations.

We now use the maximum principle of Ecker-Huisken [12, Corollary 1.1] on non-compact manifolds to show that if a smooth mean curvature flow $\{\Sigma_t\}_{t \in [-1, T]}$ satisfies that Σ_{-1} is shrinker mean convex with respect to X_0 , then this remains true for all $t \in (-1, T)$.

Proposition 3.2. *Let $\mathcal{S} = \{\Sigma_t\}_{t \in [-1, T]}$ be a smooth mean curvature flow in $\mathbb{R}^{n+1}$ with finite entropy. Suppose that Σ_{-1} satisfies*

$$(3.4) \quad S_{\Sigma_{-1}}^{X_0}(p) \geq c(1 + |\mathbf{x}(p)|^2)^{-\alpha}$$

for some $c > 0$ and $\alpha \geq 0$, and that

$$(3.5) \quad M = \sup_{t \in [-1, T]} \sup_{\Sigma_t \setminus B_R} |A_{\Sigma_t}| + |\nabla_{\Sigma_t} A_{\Sigma_t}| + |\nabla_{\Sigma_t}^2 A_{\Sigma_t}| < \infty$$

for some $R > 0$. For all $(p, t) \in \mathcal{S}$,

$$(3.6) \quad S_{\Sigma_t}^{X_0}(p) \geq c(1 + |\mathbf{x}(p)|^2 + 2n(t + 1))^{-\alpha}.$$

Proof. Let

$$(3.7) \quad \eta(\mathbf{x}, t) = 1 + |\mathbf{x}|^2 + 2n(t + 1).$$

By [12, Lemma 1.1] we have that

$$(3.8) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) \eta = 0,$$

and so

$$(3.9) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) \eta^\alpha = -\alpha(\alpha - 1) |\nabla_{\Sigma_t} \log \eta|^2 \eta^\alpha.$$

Define

$$(3.10) \quad u(p, t) = \eta^\alpha(\mathbf{x}(p), t) S_{\Sigma_t}^{X_0}(p) \quad \text{for } (p, t) \in \mathcal{S}.$$

Then it follows from Lemma 3.1 and (3.8) that

$$(3.11) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) u + 2\alpha \nabla_{\Sigma_t} \log \eta \cdot \nabla_{\Sigma_t} u = |A_{\Sigma_t}|^2 u + \alpha(\alpha + 1) |\nabla_{\Sigma_t} \log \eta|^2 u.$$

Notice that

$$(3.12) \quad |\nabla_{\Sigma_t} \log \eta|(p, t) = \frac{2|\mathbf{x}(p)^\top|}{1 + |\mathbf{x}(p)|^2 + 2n(t + 1)} < 2,$$

and by (3.5) and (3.11) we have that on $\mathcal{S} \setminus C_{R, T+1}(\mathbf{0}, -1)$,

$$(3.13) \quad \left| \frac{du}{dt} \right|(p, t) + \sum_{i=0}^2 |\nabla_{\Sigma_t}^i u|(p, t) \leq C(M, \alpha, X_0) (1 + |\mathbf{x}(p)|^2 + 2n(t + 1))^{\alpha+1}.$$

Invoking (3.4) and finiteness of the entropy, Theorem A.1 implies, when $R = 0$, that

$$(3.14) \quad \inf_{p \in \Sigma_t} u(p, t) \geq \inf_{p \in \Sigma_{-1}} u(p, -1) \geq c,$$

giving immediately (3.6). $\square$

We further adapt some ideas of [12] – specifically the proof of [12, Lemma 4.1] – to prove a relationship between the shrinker mean curvature and the second fundamental form for shrinker mean convex flows.

Proposition 3.3. *Let $\mathcal{S} = \{\Sigma_t\}_{t \in [-1, T]}$ be a smooth mean curvature flow in $\mathbb{R}^{n+1}$ with finite entropy. Suppose that Σ_{-1} satisfies*

$$(3.15) \quad S_{\Sigma_{-1}}^{X_0}(p) \geq c (1 + |\mathbf{x}(p)|^2)^{-\alpha}$$

for some $c > 0$ and $\alpha \geq 0$, and that

$$(3.16) \quad \tilde{M} = \max \left\{ \sup_{t \in [-1, T]} \sup_{\Sigma_t \setminus B_R} |A_{\Sigma_t}| + |\nabla_{\Sigma_t} A_{\Sigma_t}| + |\nabla_{\Sigma_t}^2 A_{\Sigma_t}|, \sup_{\Sigma_{-1}} |A_{\Sigma_{-1}}| \right\} < \infty$$

for some $R > 0$. Then for all $(p, t) \in \mathcal{S}$,

$$(3.17) \quad |A_{\Sigma_t}(p)| \leq c^{-1} \tilde{M} (1 + |\mathbf{x}(p)|^2 + R^2 + 2n(t + 1))^\alpha S_{\Sigma_t}^{X_0}(p).$$

Proof. First observe that, by Proposition 3.2, for all $(p, t) \in \mathcal{S}$,

$$(3.18) \quad S_{\Sigma_t}^{X_0}(p) \geq c (1 + |\mathbf{x}(p)|^2 + 2n(t+1))^{-\alpha}.$$

Thus, in view of (3.16), it suffices to prove (3.17) on $\mathcal{S} \cap C_{R,T+1}^-(\mathbf{0}, T)$.

Let

$$(3.19) \quad u(p, t) = |A_{\Sigma_t}(p)|^2 v^2(p, t) = |A_{\Sigma_t}(p)|^2 |S_{\Sigma_t}^{X_0}(p)|^{-2}.$$

By [11, Appendix B, (B.9)],

$$(3.20) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) |A_{\Sigma_t}|^2 \leq -2 |\nabla_{\Sigma_t} |A_{\Sigma_t}||^2 + 2 |A_{\Sigma_t}|^4.$$

And by Lemma 3.1,

$$(3.21) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) v^2 = -2 |A_{\Sigma_t}|^2 v^2 - 6 |\nabla_{\Sigma_t} v|^2$$

Thus, we compute the evolution equation of u :

$$(3.22) \quad \begin{aligned} \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) u &= |A_{\Sigma_t}|^2 \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) v^2 - 2 \nabla_{\Sigma_t} v^2 \cdot \nabla_{\Sigma_t} |A_{\Sigma_t}|^2 \\ &\quad + v^2 \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) |A_{\Sigma_t}|^2 \\ &\leq -6 |A_{\Sigma_t}|^2 |\nabla_{\Sigma_t} v|^2 - 2 \nabla_{\Sigma_t} v^2 \cdot \nabla_{\Sigma_t} |A_{\Sigma_t}|^2 \\ &\quad - 2 |\nabla_{\Sigma_t} |A_{\Sigma_t}||^2 v^2. \end{aligned}$$

Using Young's inequality, we obtain the estimate:

$$(3.23) \quad \begin{aligned} -2 \nabla_{\Sigma_t} v^2 \cdot \nabla_{\Sigma_t} |A_{\Sigma_t}|^2 &= - \frac{\nabla_{\Sigma_t} v^2}{v^2} \cdot \nabla_{\Sigma_t} (|A_{\Sigma_t}|^2 v^2) + \frac{|A_{\Sigma_t}|^2 |\nabla_{\Sigma_t} v^2|^2}{v^2} \\ &\quad - \nabla_{\Sigma_t} v^2 \cdot \nabla_{\Sigma_t} |A_{\Sigma_t}|^2 \\ &= - \frac{\nabla_{\Sigma_t} v^2}{v^2} \cdot \nabla_{\Sigma_t} u + 4 |A_{\Sigma_t}|^2 |\nabla_{\Sigma_t} v|^2 \\ &\quad - 4v |A_{\Sigma_t}| |\nabla_{\Sigma_t} v \cdot \nabla_{\Sigma_t} |A_{\Sigma_t}|| \\ &\leq - \frac{\nabla_{\Sigma_t} v^2}{v^2} \cdot \nabla_{\Sigma_t} u + 6 |A_{\Sigma_t}|^2 |\nabla_{\Sigma_t} v|^2 \\ &\quad + 2v^2 |\nabla_{\Sigma_t} |A_{\Sigma_t}||^2. \end{aligned}$$

Hence,

$$(3.24) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) u \leq - \frac{\nabla_{\Sigma_t} v^2}{v^2} \cdot \nabla_{\Sigma_t} u = -2 \nabla_{\Sigma_t} \log v \cdot \nabla_{\Sigma_t} u.$$

Observe that the hypothesis (3.16) and (3.18) imply that

$$(3.25) \quad \sup_{\mathcal{S} \cap \partial_P C_{R,t+1}^-(\mathbf{0}, t)} u \leq c^{-2} \tilde{M}^2 (1 + R^2 + 2n(t+1))^{2\alpha}.$$

Therefore the standard parabolic maximum principle implies that

$$(3.26) \quad \sup_{\mathcal{S} \cap C_{R,t+1}^-(\mathbf{0}, t)} u \leq c^{-2} \tilde{M}^2 (1 + R^2 + 2n(t+1))^{2\alpha},$$

which proves the result. $\square$

4. SELF-SHRINKERS OF SMALL ENTROPY ASYMPTOTIC TO REGULAR CONES

We define $\mathcal{ACS}_n$ to be the space of connected asymptotically conical self-shrinkers in $\mathbb{R}^{n+1}$. Observe that the hyperplanes through the origin are contained in $\mathcal{ACS}_n$. We denote by $\mathcal{ACS}_n^*$ the subspace of non-flat elements of $\mathcal{ACS}_n$. Furthermore, given a $\lambda \geq 1$, we let $\mathcal{ACS}_n[\lambda]$ be the set of elements of $\mathcal{ACS}_n$ with Gaussian surface area (and hence entropy) less than or equal to λ and define $\mathcal{ACS}_n^*[\lambda]$ likewise.

The goal of this section is to prove the long-time existence of a smooth mean curvature flow starting from a perturbation of any element of $\mathcal{ACS}_n^*[\lambda_{n-1}]$. First we need the following properties of the lowest eigenfunction of the stability operator.

Proposition 4.1. *Given $\Sigma \in \mathcal{ACS}_n^*$, there is a $\mu = \mu(\Sigma) < -1$ and a unique positive smooth function f on Σ which satisfies*

$$(4.1) \quad L_\Sigma f = \Delta_\Sigma f - \frac{\mathbf{x}}{2} \cdot \nabla_\Sigma f + |A_\Sigma|^2 f + \frac{1}{2} f = -\mu f \quad \text{with} \quad \int_\Sigma f^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n = 1.$$

Moreover, for each $\beta > 0$, there are constants $C_0, \dots, C_m, \dots > 1$ depending on Σ and β so that

$$(4.2) \quad C_0^{-1} (1 + |\mathbf{x}(p)|^2)^{\frac{1}{2} + \mu - \beta} < f(p) < C_0 (1 + |\mathbf{x}(p)|^2)^{\frac{1}{2} + \mu + \beta}, \quad \text{and}$$

$$(4.3) \quad |\nabla_\Sigma^m f(p)| < C_m (1 + |\mathbf{x}(p)|^2)^{\frac{1}{2} + \mu + \beta - \frac{m}{2}} \quad \text{for all } m \geq 1.$$

Proof. Given functions ϕ, ψ on Σ , we define

$$(4.4) \quad (\phi, \psi)_0 = \int_\Sigma \phi \psi e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n, \quad (\phi, \psi)_1 = (\phi, \psi)_0 + (\nabla_\Sigma \phi, \nabla_\Sigma \psi)_0,$$

and $\|\phi\|_i^2 = (\phi, \phi)_i$ for $i = 0, 1$. Let $L_w^2(\Sigma)$ be the Hilbert space of functions on Σ with finite $\|\cdot\|_0$ -norm and $H_w^1(\Sigma)$ be the Hilbert space given by completing $C_c^\infty(\Sigma)$ using $\|\cdot\|_1$. Observe that $H_w^1(\Sigma)$ may be naturally identified with a subspace of $L_w^2(\Sigma)$. In fact, this embedding is compact – see Appendix B or [7, Theorem 3].

Denote by B_R^Σ the intrinsic geodesic open ball in Σ centered at a fixed point $p \in \Sigma$ with radius R . Note that $\bigcup_{R>0} B_R^\Sigma = \Sigma$ as Σ is connected. As observed in [10, Remark 5.17], there is a $\mu_R = \mu_R(\Sigma)$ and a unique positive function $f_R \in H_w^1(B_R^\Sigma)$ so that $L_\Sigma f_R + \mu_R f_R = 0$ with $\|f_R\|_0^2 = 1$. Moreover, μ_R is characterized by

$$(4.5) \quad \mu_R = \inf_\phi \frac{\int_\Sigma \left(|\nabla_\Sigma \phi|^2 - |A_\Sigma|^2 \phi^2 - \frac{1}{2} \phi^2 \right) e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n}{\int_\Sigma \phi^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n},$$

where the infimum is taken over all smooth non-zero functions compactly supported in B_R^Σ . As $\Sigma \in \mathcal{ACS}_n^*$, it follows from [25, Lemma 2.1] that

$$(4.6) \quad (1 + |\mathbf{x}(p)|^2) |A_\Sigma(p)|^2 = O(1).$$

Obviously, μ_R is non-increasing in R . Thus, $\mu_R \rightarrow \mu$ as $R \rightarrow \infty$ for some $\mu > -\infty$. This further implies that $\|f_R\|_1^2$ is uniformly bounded. Hence, there is a sequence $R_j \rightarrow \infty$ so that f_{R_j} converges to a function f on Σ weakly in H_w^1 and strongly in L_w^2 . Clearly, $f \in H_w^1(\Sigma)$ satisfying that $L_\Sigma f + \mu f = 0$ with $\|f\|_0^2 = 1$. The smoothness of f follows from a standard elliptic regularity theory. As each f_{R_j} is non-negative, so is f and, in fact, $f > 0$ on Σ by the Harnack inequality. Finally, it follows from [10, Lemma 9.25] that such an f is unique.

As Σ is non-flat, the classification of mean convex self-shrinkers [10, Theorem 0.17] implies that H_Σ must change sign. Hence, [10, Theorem 9.36] implies that $\mu < -1$. For $\beta > 0$, define

$$(4.7) \quad \bar{g}(\mathbf{x}) = |\mathbf{x}|^{1+2\mu+2\beta} \quad \text{and} \quad \underline{g}(\mathbf{x}) = |\mathbf{x}|^{1+2\mu-2\beta}.$$

Then, invoking [10, Lemma 3.20] and (4.6), a direct computation gives

$$(4.8) \quad L_\Sigma \bar{g} + \mu \bar{g} < 0 \quad \text{and} \quad L_\Sigma \underline{g} + \mu \underline{g} > 0 \quad \text{on } \Sigma \setminus \bar{B}_\mathcal{R},$$

for some $\mathcal{R} = \mathcal{R}(\beta, \Sigma) > 1$ sufficiently large. Next we choose $C > 1$ so that

$$(4.9) \quad C^{-1} \underline{g} < f < C \bar{g} \quad \text{on } \partial(\Sigma \setminus B_\mathcal{R}).$$

Consider the Dirichlet problem

$$(4.10) \quad \begin{cases} L_\Sigma g + \mu g = 0 & \text{on } \Sigma \cap (B_i \setminus \bar{B}_\mathcal{R}), \\ g = f & \text{on } \partial(\Sigma \setminus B_\mathcal{R}), \\ g = C \bar{g} & \text{on } \partial(\Sigma \setminus B_i). \end{cases}$$

Note that the zeroth order term of the differential equation in (4.10) has a negative sign. Thus, it follows from [14, Theorems 8.3 and 8.13] and the maximum principle that for each $i > \mathcal{R}$, the problem (4.10) has a unique smooth solution $g = g_i$ bounded between $C^{-1} \underline{g}$ and $C \bar{g}$. Passing $i \rightarrow \infty$, by the Arzela-Ascoli theorem, g_i converges in $C_{loc}^\infty(\Sigma \setminus B_\mathcal{R})$ to some function $\tilde{g}$ so that $L_\Sigma \tilde{g} + \mu \tilde{g} = 0$ and $\tilde{g} = f$ on $\partial(\Sigma \setminus B_\mathcal{R})$. Moreover,

$$(4.11) \quad C^{-1} |\mathbf{x}(p)|^{1+2\mu-2\beta} < \tilde{g}(p) < C |\mathbf{x}(p)|^{1+2\mu+2\beta}.$$

It follows from (4.6) and elliptic Schauder estimates that $|\nabla_\Sigma \tilde{g}|$ grows at most polynomially. This together with the Euclidean volume growth of Σ implies that

$$(4.12) \quad \int_{\Sigma \setminus B_\mathcal{R}} |\nabla_\Sigma \tilde{g}|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n < \infty.$$

To conclude the proof, it suffices to show that $h = f - \tilde{g} = 0$ on $\Sigma \setminus B_\mathcal{R}$. Let ϕ_i be a cut-off function on $\mathbb{R}^{n+1}$ so that $\phi_i = 1$ in B_i , $\phi_i = 0$ outside B_{2i} , and $|D\phi_i| < 2/i$. Since $L_\Sigma h + \mu h = 0$ with $h = 0$ on $\partial(\Sigma \setminus B_\mathcal{R})$, multiplying both sides of the equation by $h \phi_i^2 e^{-|\mathbf{x}|^2/4}$ and integration by parts give that

$$(4.13) \quad \begin{aligned} \int_{\Sigma \setminus B_\mathcal{R}} |\nabla_\Sigma h|^2 \phi_i^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n - \int_{\Sigma \setminus B_\mathcal{R}} \left(|A_\Sigma|^2 + \frac{1}{2} + \mu \right) h^2 \phi_i^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n \\ + \int_{\Sigma \cap (B_{2i} \setminus B_i)} 2h \phi_i \nabla_\Sigma \phi_i \cdot \nabla_\Sigma h e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n = 0. \end{aligned}$$

Thus, sending $i \rightarrow \infty$ and invoking that $\mu < -1$, the monotone convergence theorem together with (4.11) and (4.12) implies that $h = 0$ on $\Sigma \setminus B_\mathcal{R}$ and so (4.2) follows.

Finally, letting $\Sigma_t = \sqrt{-t} \Sigma$, we define

$$(4.14) \quad \tilde{f}(q, t) = (-t)^{\mu+\frac{1}{2}} f\left(\frac{q}{\sqrt{-t}}\right) \quad \text{for } t < 0 \text{ and } q \in \Sigma_t.$$

Then (4.1) implies that

$$(4.15) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) \tilde{f} = |A_{\Sigma_t}|^2 \tilde{f}.$$

And (4.2) gives that, if $t \in [-1, 0)$ and $q \in \Sigma_t \cap (B_2 \setminus B_1)$, then

$$(4.16) \quad 0 < \tilde{f}(q, t) < C'(\Sigma, \mu, \beta)(-t)^{-\beta}.$$

Recall that $\Sigma_t \rightarrow \mathcal{C}(\Sigma)$ in $C_{loc}^\infty(\mathbb{R}^{n+1} \setminus \{0\})$ as $t \rightarrow 0$. Thus, it follows from standard parabolic regularity theory that for each integer $m \geq 1$, there is a $C'' = C''(\Sigma, C', m)$ so that if $p \in \Sigma \setminus B_R$, then

$$(4.17) \quad |\nabla_\Sigma^m f(p)| \leq C'' |\mathbf{x}(p)|^{1+2\mu+2\beta-m},$$

proving (4.3). $\square$

We next use the lowest eigenfunction of the stability operator to perturb any element of $\mathcal{ACS}_n^*$ in order to strictly decrease its entropy.

Proposition 4.2. *Let $\Sigma \in \mathcal{ACS}_n^*$ and let C_0 , μ , and f be as in Proposition 4.1. There is an $\epsilon_0 = \epsilon_0(\Sigma) \in (0, 1)$ and $K = K(\Sigma) > 1$ so that if $|\epsilon| < \epsilon_0$, then:*

(1) *the normal graph over Σ given by*

$$(4.18) \quad \Gamma^\epsilon = \{\mathbf{y} \in \mathbb{R}^{n+1} : \mathbf{y} = \mathbf{x}(p) + \epsilon f(p) \mathbf{n}_\Sigma(p), p \in \Sigma\}$$

is a smooth hypersurface;

(2) *for all $R > 1$ we have that $\Gamma^\epsilon \setminus B_{KR} \subset \mathcal{T}_{1/R}(\mathcal{C}(\Sigma))$;*

(3) *Γ^ϵ is asymptotically conical and, in particular, given $\delta > 0$ there is a $\kappa \in (0, 1)$ and $\mathcal{R} > 1$ depending only on δ and Σ so that if $p \in \Sigma \setminus B_{\mathcal{R}}$ and $r = \kappa |\mathbf{x}(p)|$, then $\Gamma^\epsilon \cap B_r(p)$ can be written as a connected graph of a function w over $T_p \Sigma$ with $|Dw| \leq \delta$;*

(4) *by a suitable choice of the normal to Γ^ϵ ,*

$$(4.19) \quad S_{\Gamma^\epsilon}^{O, -1}(p) \geq -\mu |\epsilon| C_0^{-1} (1 + |\mathbf{x}(p)|^2)^\mu \geq 0;$$

(5) *if $\epsilon \neq 0$, then $\lambda[\Gamma^\epsilon] < \lambda[\Sigma]$.*

Proof. By Proposition 4.1, $\mu < -1$, and so if $\beta = \frac{1}{2} \min\{1, -\mu - 1\}$, then $0 < \beta \leq 1/2$ and $\mu + \beta < -1$. As μ depends only on Σ , so does β . Thus the constants C_0, C_1 in Proposition 4.1 depend only on Σ as well. Now define the mapping

$$(4.20) \quad \Xi_\epsilon : \Sigma \rightarrow \Gamma^\epsilon \subset \mathbb{R}^{n+1}, \quad \Xi_\epsilon(p) = \mathbf{x}(p) + \epsilon f(p) \mathbf{n}_\Sigma(p).$$

As $\Sigma \in \mathcal{ACS}_n^*$, in view of [25, Lemma 2.2], there is a $\rho > 0$ so that $\mathcal{T}_\rho(\Sigma)$ is a regular tubular neighborhood. By Proposition 4.1, f is smooth and uniformly bounded by C_0 and so, if $|\epsilon| < \rho/C_0$, then $\Xi_\epsilon(p) = \Psi_\rho(p, \epsilon f(p))$ and so Γ^ϵ is the image under the diffeomorphism Ψ_ρ of the graph of ϵf in $\Sigma \times (-\rho, \rho)$. In particular Γ^ϵ is a hypersurface, proving Item (1).

As $\mu + \beta < -1$, Item (2) follows directly from (4.2) for sufficiently large $K = K(\Sigma)$.

Let $\delta > 0$ be any given small constant and $\mathcal{R} = \mathcal{R}(\delta, \Sigma)$ be a sufficiently large constant which may change among lines. By [25, Lemma 2.2], there is a $\kappa = \kappa(\delta, \Sigma) \in (0, 1)$ so that if $p \in \Sigma \setminus B_{\mathcal{R}}$ and $r = \kappa |\mathbf{x}(p)|$, then $\Sigma \cap B_{2r}(p)$ is given by a connected graph of a function over $T_p \Sigma$ with gradient bounded by $\delta/2$. As β was chosen so $\beta + \mu < -1$, (4.2) and (4.3) together with [25, Lemma 2.1], and a direct computation (see the proof of [25, Lemma 2.4]) imply that if $q \in \Sigma \cap B_{2r}(p)$, then

$$(4.21) \quad |\mathbf{n}_{\Gamma^\epsilon}(\Xi_\epsilon(q)) \cdot \mathbf{n}_\Sigma(q)| > 1 - C\epsilon (1 + |\mathbf{x}(p)|)^{-2},$$

where $C > 0$ depends only on C_0, C_1 and Σ . For $|\epsilon| < \rho/C_0$, $|\Xi_\epsilon(p) - p| < \rho$ and so

$$(4.22) \quad \Gamma^\epsilon \cap B_r(p) \subset \Xi_\epsilon(\Sigma \cap B_{2r}(p)).$$

Combining (4.21) and (4.22), if $p \in \Sigma \setminus B_{\mathcal{R}}$, we conclude that $\Gamma^\epsilon \cap B_r(p)$ can be written as a connected graph of a function over $T_p \Sigma$ with gradient bounded by δ . In view of (4.2)

and (4.3), it follows from Item (2) that Γ^ϵ is smoothly asymptotic to $\mathcal{C}(\Sigma)$. This completes the proof of Item (3).

As computed in the proof of [25, Lemma 2.4], it follows from an appropriate choice of unit normal to Γ^ϵ and (4.1) that

$$(4.23) \quad S_{\Gamma^\epsilon}^{O,-1} = 2|\epsilon|L_\Sigma f + Q_\Sigma(\epsilon f, \epsilon \nabla_\Sigma f, \epsilon \nabla_\Sigma^2 f) = -2|\epsilon|\mu f + Q_\Sigma(\epsilon f, \epsilon \nabla_\Sigma f, \epsilon \nabla_\Sigma^2 f).$$

Where Q_Σ is a polynomial all of whose terms are of order at least 2 and which has uniformly bounded coefficients. This, together with (4.2), (4.3) and the fact that $\beta \leq 1/2$, implies Item (4) for all $|\epsilon|$ sufficiently small.

Finally, Item (5) can be shown by modifying the arguments in the proof of [10, Theorem 0.15]. We leave the technical details to Appendix C. $\square$

We next use sphere barriers to show that smooth mean curvature flows which initially decay to a regular cone must continue to decay to the same cone for all positive time.

Lemma 4.3. *Let $\{\Gamma_t\}_{t \in [-1, T]}$ be a smooth mean curvature flow in $\mathbb{R}^{n+1}$. Suppose that there is a regular cone, $\mathcal{C}$, and a $K > 1$ so that $\Gamma_{-1} \setminus B_{KR} \subset \mathcal{T}_{1/R}(\mathcal{C})$ for all $R > 1$. Then there is a constant $K' = K'(\mathcal{C}, K) > 1$ so that for all $R > 1$ and $t \in [-1, T]$,*

$$(4.24) \quad \Gamma_t \setminus B_{K'(R + \sqrt{2n(t+1)})} \subset \mathcal{T}_{K'(1+2n(t+1))R^{-1}}(\mathcal{C}).$$

Proof. For $\mathbf{x} \in \mathbb{R}^{n+1}$, let $\varrho(\mathbf{x}) = \inf \{ \rho \geq 0 : B_\rho(\mathbf{x}) \cap (B_{KR} \cup \mathcal{T}_{1/R}(\mathcal{C})) \neq \emptyset \}$. Define an open subset of $\mathbb{R}^{n+1}$ by

$$(4.25) \quad U = \bigcup_{\mathbf{x} \in \mathbb{R}^{n+1}} B_{\varrho(\mathbf{x})}(\mathbf{x}).$$

Clearly, $U \cap \Gamma_{-1} = \emptyset$. For $t \geq -1$, set

$$(4.26) \quad \varrho(\mathbf{x}, t) = \begin{cases} \sqrt{\varrho(\mathbf{x})^2 - 2n(t+1)} & \varrho(\mathbf{x})^2 \geq 2n(t+1) \\ 0 & \varrho(\mathbf{x})^2 < 2n(t+1), \end{cases}$$

$$(4.27) \quad U_t = \bigcup_{\mathbf{x} \in \mathbb{R}^{n+1}} B_{\varrho(\mathbf{x}, t)}(\mathbf{x}).$$

By the avoidance principle for mean curvature flow, $U_t \cap \Gamma_t = \emptyset$ for all $t \in [-1, T]$.

Let $d_{\mathcal{C}}$ denote the distance to the cone $\mathcal{C}$. Observe that

$$(4.28) \quad \varrho(\mathbf{x}) = \max \{ \min \{ |\mathbf{x}| - KR, d_{\mathcal{C}}(\mathbf{x}) - R^{-1} \}, 0 \}$$

and so the set $U_t^+ = \{ \mathbf{x} \in \mathbb{R}^{n+1} : \varrho(\mathbf{x}, t) > 0 \}$ satisfies

$$(4.29) \quad U_t^+ = \left\{ \mathbf{x} \in \mathbb{R}^{n+1} : |\mathbf{x}| > KR + \sqrt{2n(t+1)}, d_{\mathcal{C}}(\mathbf{x}) > R^{-1} + \sqrt{2n(t+1)} \right\}.$$

Clearly, $U_t^+ \subset U_t$ and, in general, is a proper subset. Hence,

$$(4.30) \quad \Gamma_t \setminus B_{KR + \sqrt{2n(t+1)}} \subset \mathcal{T}_{R^{-1} + \sqrt{2n(t+1)}}(\mathcal{C}).$$

To conclude, pick a unit normal $\mathbf{n}_{\mathcal{C}}$ to $\mathcal{C} \setminus \{0\}$ so there is a smooth map

$$(4.31) \quad \Psi : \mathcal{C} \setminus \{0\} \times \mathbb{R} \rightarrow \mathbb{R}^{n+1}, \quad \Psi(p, l) = \mathbf{x}(p) + l\mathbf{n}_{\mathcal{C}}(p).$$

As $\mathcal{L}(\mathcal{C})$, the link of $\mathcal{C}$, is compact and smooth, there is a $\vartheta \in (0, 1/2)$ so that if

$$(4.32) \quad N = \{ (p, l) \in \mathcal{C} \times \mathbb{R} : |l| < 2\vartheta |\mathbf{x}(p)| \},$$

then $\Psi|_N$ is a diffeomorphism onto its image.

Fix $t > -1$ and consider the part of U_t^c far from the origin. That is, set $K' = 1 + 2K\vartheta^{-1} > 4K$ and

$$(4.33) \quad V_t = U_t^c \cap \left\{ \mathbf{x} \in \mathbb{R}^{n+1} : |\mathbf{x}| \geq K' \left(R + \sqrt{2n(t+1)} \right) \right\}.$$

If $\mathbf{y} \in V_t$, then, by the definition of U_t and the fact that $|\mathbf{y}| \geq K'R > KR$, $d_{\mathcal{C}}(\mathbf{y}) \leq R^{-1} + \sqrt{2n(t+1)}$. Hence, there is a point $p \in \mathcal{C}$ so $d_{\mathcal{C}}(\mathbf{y}) = |\mathbf{y} - \mathbf{x}(p)| \leq R^{-1} + \sqrt{2n(t+1)}$. This implies that

$$(4.34) \quad \begin{aligned} |\mathbf{x}(p)| &\geq K' \left(R + \sqrt{2n(t+1)} \right) - \sqrt{2n(t+1)} - R^{-1} \\ &\geq (K' - 1) \left(R + \sqrt{2n(t+1)} \right) \\ &= 2K\vartheta^{-1} \left(R + \sqrt{2n(t+1)} \right), \end{aligned}$$

where the second inequality used that $R > 1$ and so $R > R^{-1}$. Hence,

$$(4.35) \quad \frac{|\mathbf{y} - \mathbf{x}(p)|}{|\mathbf{x}(p)|} \leq \frac{R^{-1} + \sqrt{2n(t+1)}}{2K\vartheta^{-1} \left(R + \sqrt{2n(t+1)} \right)} < \vartheta,$$

where we used that $R > R^{-1}$ and that $K > 1$.

Thus, $\mathbf{y} \in \Psi(N)$ and so, up to reversing the orientation of $\mathcal{C}$, $\mathbf{y} = \Psi(p, d_{\mathcal{C}}(\mathbf{y})) = \mathbf{x}(p) + d_{\mathcal{C}}(\mathbf{y})\mathbf{n}_{\mathcal{C}}(p)$. Set $\mathbf{y}_0 = \mathbf{x}(p) + \vartheta|\mathbf{x}(p)|\mathbf{n}_{\mathcal{C}}(p)$. Observe that

$$(4.36) \quad |\mathbf{y}_0| - (\vartheta|\mathbf{x}(p)| - R^{-1}) = \left(\sqrt{1 + \vartheta^2} - \vartheta \right) |\mathbf{x}(p)| + R^{-1} > KR,$$

where we used that, by (4.34), $|\mathbf{x}(p)| > 4KR$ and $\sqrt{1 + \vartheta^2} - \vartheta > 1/3$. Thus,

$$(4.37) \quad B_{\vartheta|\mathbf{x}(p)| - R^{-1}}(\mathbf{y}_0) \cap B_{KR} = \emptyset \quad \text{and so} \quad \varrho(\mathbf{y}_0) = \vartheta|\mathbf{x}(p)| - R^{-1}.$$

As $\mathbf{y} \notin U_t$, we have $\mathbf{y} \notin B_{\rho(\mathbf{y}_0, t)}(\mathbf{y}_0)$. That is, $\rho(\mathbf{y}_0, t) \leq |\mathbf{y} - \mathbf{y}_0| = \vartheta|\mathbf{x}(p)| - d_{\mathcal{C}}(\mathbf{y})$. Hence,

$$(4.38) \quad \begin{aligned} d_{\mathcal{C}}(\mathbf{y}) &\leq \vartheta|\mathbf{x}(p)| - \sqrt{(\vartheta|\mathbf{x}(p)| - R^{-1})^2 - 2n(t+1)} \\ &= \frac{\vartheta^2|\mathbf{x}(p)|^2 - (\vartheta|\mathbf{x}(p)| - R^{-1})^2 + 2n(t+1)}{\vartheta|\mathbf{x}(p)| + \sqrt{(\vartheta|\mathbf{x}(p)| - R^{-1})^2 - 2n(t+1)}} \\ &< \frac{2R^{-1}\vartheta|\mathbf{x}(p)| - R^{-2} + 2n(t+1)}{\vartheta|\mathbf{x}(p)|} \\ &< \frac{2}{R} + \frac{2n(t+1)}{\vartheta|\mathbf{x}(p)|} < \frac{2(1 + 2n(t+1))}{\vartheta R}, \end{aligned}$$

where we used $|\mathbf{x}(p)| > R$ for the last inequality. Therefore, as $\Gamma_t \subset U_t^c$ and $K' > 2/\vartheta$,

$$(4.39) \quad \Gamma_t \setminus B_{K' \left(R + \sqrt{2n(t+1)} \right)} \subset \mathcal{T}_{K'(1+2n(t+1))R^{-1}}(\mathcal{C}),$$

from which the result follows. $\square$

In what follows, we establish the existence and uniqueness of a smooth shrinker mean convex mean curvature flow starting from Γ^e at time -1 . Moreover, we show that each time slice of this flow is smoothly asymptotic to $\mathcal{C}(\Sigma)$ and if the flow exists smoothly beyond time 0, then the time zero slice lies strictly on one side of $\mathcal{C}(\Sigma)$.

Proposition 4.4. *Let $\Sigma \in \mathcal{ACS}_n^*$ and let Γ^ϵ be as in Proposition 4.2 for $\epsilon \neq 0$. There is a $T = T(\Gamma^\epsilon) \in (-1, \infty]$, and a unique smooth mean curvature flow $\{\Gamma_t^\epsilon\}_{t \in [-1, T)}$ with $\Gamma_{-1}^\epsilon = \Gamma^\epsilon$. Furthermore, there is an increasing function $R(t) > 1$ on $[-1, \infty)$ depending only on Γ^ϵ so that:*

- (1) *for $t \in [-1, T)$, Γ_t^ϵ is smoothly asymptotic to $\mathcal{C}(\Sigma)$ and, in particular, for $p \in \Gamma_t^\epsilon \setminus B_{R(t)}$,*

$$(4.40) \quad |\mathbf{x}(p)| |A_{\Gamma_t^\epsilon}(p)| + |\mathbf{x}(p)|^2 |\nabla_{\Gamma_t^\epsilon} A_{\Gamma_t^\epsilon}(p)| + |\mathbf{x}(p)|^3 |\nabla_{\Gamma_t^\epsilon}^2 A_{\Gamma_t^\epsilon}(p)| < \tilde{C}_0,$$

where $\tilde{C}_0$ depends only on Γ^ϵ ;

- (2) *if $T < \infty$, then*

$$(4.41) \quad \lim_{t \rightarrow T} \sup_{B_{R(t)} \cap \Gamma_t^\epsilon} |A_{\Gamma_t^\epsilon}| = \infty;$$

- (3) *for $t < \min\{0, T\}$, $\Gamma_t^\epsilon \setminus \bar{B}_{R(t)}$ is given by the normal graph of a function f_t over a region $\Omega_t \subset \Sigma_t = \sqrt{-t}\Sigma$ with $\Sigma_t \setminus B_{2R(t)} \subset \Omega_t$ which satisfies $f_{-1} = \epsilon f|_{\Omega_{-1}}$,*

$$(4.42) \quad \tilde{C}_1^{-1} |\epsilon| (1 + |\mathbf{x}(p)|)^{2\mu} < |f_t(p)| < \tilde{C}_1 (1 + |\mathbf{x}(p)|)^{-1}, \quad \text{and}$$

$$(4.43) \quad (1 + |\mathbf{x}(p)|)^2 |\nabla_{\Sigma_t} f_t(p)| + (1 + |\mathbf{x}(p)|)^3 |\nabla_{\Sigma_t}^2 f_t(p)| < \tilde{C}_1,$$

where $\tilde{C}_1$ depends only on Σ ;

- (4) *for $t < \min\{0, T\}$, $\Gamma_t^\epsilon \cap \Sigma_t = \emptyset$, and if $T > 0$, then $\Gamma_0^\epsilon \cap \mathcal{C}(\Sigma) = \emptyset$;*

- (5) *by a suitable choice of the normal to Γ^ϵ ,*

$$(4.44) \quad S_{\Gamma_t^\epsilon}^O(p) \geq -\mu |\epsilon| C_0^{-1} (1 + |\mathbf{x}(p)|^2 + 2n(t+1))^\mu > 0.$$

Proof. First, it follows from Items (2) and (3) of Proposition 4.2 that Γ^ϵ smoothly asymptotes to the same regular cone, $\mathcal{C}(\Sigma)$, as that of Σ . Thus, Γ^ϵ has uniformly bounded C^m norm for all m and $\mathcal{T}_\rho(\Gamma^\epsilon)$ is a regular tubular neighborhood of Γ^ϵ for some $\rho > 0$. Arguing as in the proof of [13, Theorem 4.2], there exists a smooth mean curvature flow starting from Γ^ϵ for small positive time. Indeed, the construction ensures that each of these time slices can be expressed as a smooth normal graph over Γ^ϵ with small L^∞ norm. Denote the maximal extension of this flow to a smooth mean curvature flow by $\{\Gamma_t^\epsilon\}_{t \in [-1, T)}$.

Invoking Item (2) of Proposition 4.2, it follows from Lemma 4.3 and [25, Lemma 2.2] that there exist increasing functions $R(t) > 1$ and $C(t) > 1$ on $[-1, \infty)$ depending only on Γ^ϵ so that for $t \in (-1, T)$ and $p \in \Gamma_t^\epsilon \setminus B_{R(t)}$, $d_\Sigma(\mathbf{x}(p)) \leq C(t) |\mathbf{x}(p)|^{-1}$, where d_Σ is the distance to Σ . Furthermore, the pseudo-locality property of mean curvature flow (cf. [18, Theorem 1.5]) together with Item (3) of Proposition 4.2 (for $\delta > 0$ sufficiently small) implies that each slice $\Gamma_\tau^\epsilon \cap B_r(q)$ is a connected graph of a function w_τ over $T_q \Sigma$ with $|Dw_\tau| \leq 2\delta$. Here, $\tau \leq t$ and

$$(4.45) \quad q \in \Sigma \text{ is such that } |\mathbf{x}(p) - \mathbf{x}(q)| = d_\Sigma(\mathbf{x}(p)) \text{ and } r = \varsigma \kappa |\mathbf{x}(q)|,$$

for some universal small constant ς . Replacing the localization function in the proof of [11, Propositions 3.21 and 3.22] with $(1 - |\mathbf{x}|^2)_+^3$, we conclude that, for $m \geq 0$,

$$(4.46) \quad \sup_{\tau \in [-1, t]} \sup_{\Gamma_\tau^\epsilon \cap B_{\frac{r}{2}}(q)} |\nabla_{\Gamma_\tau^\epsilon}^m A_{\Gamma_\tau^\epsilon}| \leq C'(m, \delta, \Gamma^\epsilon) r^{-m-1}.$$

This, together with Item (2) of Proposition 4.2 and Lemma 4.3, immediately gives Item (1) and allows us to invoke [6, Theorem 1.1] to establish that the flow $\{\Gamma_t^\epsilon\}_{t \in [-1, T)}$ is the unique smooth mean curvature flow starting from Γ^ϵ at time -1 . Hence, it remains only to verify Items (2), (3), (4) and (5).

To show Item (2), we argue by contradiction. If $T < \infty$ and

$$(4.47) \quad \sup_{t \in [-1, T)} \sup_{\Gamma_t^\epsilon} |A_{\Gamma_t^\epsilon}| < \infty,$$

then the same holds true for all higher order derivatives of the second fundamental form by [13, Theorem 3.7]. As Γ^ϵ decays smoothly to $\mathcal{C}(\Sigma)$, it follows from Item (1) that Γ_t^ϵ for $t < T$ smoothly decays to $\mathcal{C}(\Sigma)$ in a uniform manner. Thus, $\Gamma_t^\epsilon \rightarrow \Gamma_T^\epsilon$ smoothly and Γ_T^ϵ is a hypersurface in $\mathbb{R}^{n+1}$ smoothly decaying to $\mathcal{C}(\Sigma)$. By our previous discussion, we could extend smoothly the flow $\{\Gamma_t^\epsilon\}_{t \in [-1, T)}$ to a larger time interval $[-1, T')$ for some $T' > T$. This contradicts the maximality of T .

By [25, Lemma 2.2], for $\tau \leq t < \min\{0, T\}$, $p \in \Sigma_\tau$ and q and r as in (4.45), $\Sigma_\tau \cap B_r(q)$ is the connected graph of a function $\tilde{w}_\tau$ over $T_q \Sigma$ with $r^{m-1} |D^m \tilde{w}_\tau|$ uniformly bounded for $1 \leq m \leq 3$. Below, let $C'' = C''(\delta, \Sigma, \Gamma^\epsilon)$ be a sufficiently large constant which may change among lines. In view of (4.2) and (4.3), it is straightforward to check that

$$(4.48) \quad \sum_{m=0}^2 r^{-1-2\mu-2\beta+m} \|D^m w_{-1} - D^m \tilde{w}_{-1}\|_{L^\infty} \leq C''.$$

Thus, combining (4.46) and (4.48), it follows from the mean curvature flow equation that, as long as β is small enough so that $\mu + \beta < -1$, for $t < \min\{0, T\}$,

$$(4.49) \quad \sum_{m=0}^2 r^{m+1} \|D^m w_t - D^m \tilde{w}_t\|_{L^\infty} \leq C''.$$

Hence, using similar reasoning as in the proof of [25, Lemma 2.3], one verifies all of Item (3) except the lower bound on $|f_t|$, which will be a consequence of our proof of Item (4).

Next, by symmetry, it suffices to show Item (4) for $\epsilon > 0$ and so $f_{-1} = \epsilon f > 0$. Let

$$(4.50) \quad T_0 = \sup \{t_0 < \min\{0, T\} : \Gamma_t^\epsilon \cap \Sigma_t = \emptyset \text{ for all } t < t_0\}.$$

We will show that $T_0 = \min\{0, T\}$, implying the first part of Item (4), i.e., $\Gamma_t^\epsilon \cap \Sigma_t = \emptyset$ when $t < \min\{0, T\}$. First, we prove that Γ_t^ϵ remains disjoint from Σ_t for t close to -1 . By continuity, there is a $\delta' > 0$ small so that $\Gamma_t^\epsilon \cap \Sigma_t \cap \bar{B}_{2R(0)} = \emptyset$ for all $t < -1 + \delta'$. In particular, if $t < -1 + \delta'$, then f_t restricted to $\Sigma_t \cap \partial B_{2R(0)}$ is strictly positive. As both Γ_t^ϵ and Σ_t move by mean curvature, the equation for f_t is given by a perturbation of the linearized mean curvature flow equation and thus, invoking [25, Lemma 2.1] and Item (3), we get that

$$(4.51) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) f_t = \mathbf{a} \cdot \nabla_{\Sigma_t} f_t + b f_t,$$

where $\mathbf{a}$ and b are uniformly bounded and depend on Σ , f_t , $\nabla_{\Sigma_t} f_t$, and $\nabla_{\Sigma_t}^2 f_t$. The derivation of (4.51) involves lengthy but tedious computations for which we refer the reader to [21, Lemma 2.5]. Define

$$(4.52) \quad \tilde{f}_t(p) = f_t(p) \eta^{-\mu}(\mathbf{x}(p), t) = f_t(p) (1 + |\mathbf{x}(p)|^2 + 2n(t+1))^{-\mu}.$$

Then it is easy to deduce that

$$(4.53) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) \tilde{f}_t = \tilde{\mathbf{a}} \cdot \nabla_{\Sigma_t} \tilde{f}_t + \tilde{b} \tilde{f}_t,$$

where

$$(4.54) \quad \tilde{\mathbf{a}} = \mathbf{a} + 2\mu \nabla_{\Sigma_t} \log \eta \quad \text{and} \quad \tilde{b} = b + \mu(\mu - 1) |\nabla_{\Sigma_t} \log \eta|^2 + \mu \mathbf{a} \cdot \nabla_{\Sigma_t} \log \eta.$$

Thus, together with (4.2) (choosing $\beta = 1/2$), it follows from Theorem A.1 that, for $t < -1 + \delta'$, $\tilde{f}_t$ in $\Sigma_t \setminus B_{2R(0)}$ is bounded from below by a positive constant. In particular, this implies that $T_0 \geq -1 + \delta'$.

Assume that $T_0 < \min\{0, T\}$. Invoking Item (3) and [25, Lemma 2.1] again, it follows from the parabolic Harnack inequality [19, Corollary 7.42] that $\tilde{f}_t$ restricted on $\Sigma_t \setminus B_{4R(0)}$ for $t \leq T_0$ is also bounded from below by a positive constant. Thus, by the parabolic maximum principle on bounded domains, we conclude that $\Gamma_{T_0}^\epsilon \cap \Sigma_{T_0} = \emptyset$. Hence, we repeat the arguments in the previous paragraph to see that $\Gamma_t^\epsilon \cap \Sigma_t = \emptyset$ for $t \in [T_0, T'_0]$ and $T'_0 > T_0$. This contradicts the definition of T_0 . Therefore, $T_0 = \min\{0, T\}$, i.e., $\Gamma_t^\epsilon \cap \Sigma_t = \emptyset$ for $t < \min\{0, T\}$.

To conclude, we note that, as $\Sigma_t \rightarrow \mathcal{C}(\Sigma)$ in $C_{loc}^\infty(\mathbb{R}^{n+1} \setminus \{0\})$ as $t \rightarrow 0$, we must have $\Gamma_0^\epsilon \cap \mathcal{C}(\Sigma) \subset \{0\}$ by the previous discussion and parabolic maximum principle on bounded domains. However, if $\Gamma_0^\epsilon \cap \mathcal{C}(\Sigma) = \{0\}$, then, as $\{\Gamma_t^\epsilon\}_{t \in [-1, 0]}$ is smooth, its parabolic blow-up at O would be a static hyperplane and so Σ would lie in a half-space whose boundary is this hyperplane. By [10, Lemma 3.25] and the fact that Σ has polynomial volume growth, this could occur only if Σ was the hyperplane – see also [5, Theorem 1.1]. This contradicts $\Sigma \in \mathcal{ACS}_n^*$ and shows Item (4). Finally, Item (5) follows from Proposition 3.2, Item (1) and Item (4) of Proposition 4.2. $\square$

Following the arguments in [9], we show that, for elements of $\mathcal{ACS}_n^*$ with small entropy, the flow constructed in Proposition 4.4 exists for long time. A crucial ingredient in our proof is the fact that shrinker mean convexity is preserved under mean curvature flow – i.e., Item (5) of Proposition 4.4.

Proposition 4.5. *If $\Sigma \in \mathcal{ACS}_n^*[\lambda_{n-1}]$ and, for $\epsilon \neq 0$, $\{\Gamma_t^\epsilon\}_{t \in [-1, T]}$ is given by Proposition 4.4, then $T = \infty$.*

Proof. We argue by contradiction. If $T < \infty$, then Item (5) of Proposition 4.2, Item (2) of Proposition 4.4 and Brakke's local regularity theorem [3] or [26] imply that there is an $\mathbf{x}_0 \in \bar{B}_{R(T)}$ so that the corresponding rescaled flow about $X_0 = (\mathbf{x}_0, T)$,

$$(4.55) \quad \tilde{\Gamma}_s^\epsilon = (T - t)^{-\frac{1}{2}} (\Gamma_t^\epsilon - \mathbf{x}_0), \quad s = -\log(T - t),$$

satisfies that for some sequence $s_i \rightarrow \infty$, $\tilde{\Gamma}_{s_i}^\epsilon$ converges to a multiplicity one F -stationary varifold $\tilde{\Sigma}$ with $1 < \lambda[\tilde{\Sigma}] < \lambda_{n-1}$ and

$$(4.56) \quad \int_{\tilde{\Gamma}_{s_i}^\epsilon} \left| \mathbf{H}_{\tilde{\Gamma}_{s_i}^\epsilon} + \frac{\mathbf{x}^\perp}{2} \right|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n \rightarrow 0.$$

Invoking Items (3) and (4) of Proposition 4.2 and Item (1) of Proposition 4.4, it follows from Proposition 3.3 that on $\tilde{\Gamma}_{s_i}^\epsilon \cap B_{R_i}$ for $R_i = R(T)e^{s_i/2}$,

$$(4.57) \quad \left| A_{\tilde{\Gamma}_{s_i}^\epsilon} \right| < C(\Sigma, \Gamma^\epsilon, T) \left\{ 2(e^{-s_i} - T) H_{\tilde{\Gamma}_{s_i}^\epsilon} - e^{-\frac{s_i}{2}} \left(\mathbf{x}_0 + e^{-\frac{s_i}{2}} \mathbf{x} \right) \cdot \mathbf{n}_{\tilde{\Gamma}_{s_i}^\epsilon} \right\}.$$

Passing $s_i \rightarrow \infty$, this together with Brakke's local regularity theorem implies that

$$(4.58) \quad |A_{\tilde{\Sigma}}| \leq -2CTH_{\tilde{\Sigma}} \quad \text{on the regular set } \text{Reg}(\tilde{\Sigma}).$$

If $n \geq 3$, then $\lambda_{n-1} < 3/2$ and so $\tilde{\Sigma}$ is smoothly embedded by [9, Proposition 5.1]. If $n = 2$, it follows from (4.56) and (4.57) that

$$(4.59) \quad \int_{\tilde{\Gamma}_{s_i}^\epsilon \cap B_R} \left| A_{\tilde{\Gamma}_{s_i}^\epsilon} \right|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^2 < C'(\lambda_1, R, T, C).$$

Thus, in view of [22, Lemma 4], the singular set $\text{Sing}(\tilde{\Sigma})$ is discrete. Furthermore, as $\tilde{\Sigma}$ is of multiplicity one, it follows from Allard's regularity theorem that $\text{Sing}(\tilde{\Sigma}) = \emptyset$.

If $T = 0$, then $\tilde{\Sigma}$ is flat and of multiplicity one and hence, $\lambda[\tilde{\Sigma}] = 1$, contradicting $\lambda[\tilde{\Sigma}] > 1$. If $T \neq 0$, then $H_{\tilde{\Sigma}}$ does not change sign and, as $F[\tilde{\Sigma}] < \lambda_{n-1}$, it follows from [10, Theorem 0.14] that $\tilde{\Sigma}$ is the self-shrinking sphere or a hyperplane. As $\tilde{\Gamma}_{s_i}^\epsilon$ is connected and non-compact, $\tilde{\Sigma}$ must be also non-compact and so $\tilde{\Sigma}$ cannot be a sphere. Hence, $\tilde{\Sigma}$ must be a hyperplane, yielding a contradiction as in the previous case. $\square$

We will need the following straightforward consequence of Propositions 4.4 and 4.5 in the next section.

Theorem 4.6. *Let $\Sigma \in \mathcal{ACS}_n^*[\lambda_{n-1}]$ and let Γ^ϵ be as in Proposition 4.2 for $\epsilon \neq 0$. There is a unique smooth mean curvature flow $\{\Gamma_t^\epsilon\}_{t \in [-1, 0]}$ with $\Gamma_{-1}^\epsilon = \Gamma^\epsilon$. Moreover, Γ_0^ϵ is a connected hypersurface which is smoothly asymptotic to, but disjoint from, $\mathcal{C}(\Sigma)$, and which satisfies $\mathbf{x} \cdot \mathbf{n}_{\Gamma_0^\epsilon} > 0$.*

5. TOPOLOGY OF NON-COMPACT SHRINKERS OF SMALL ENTROPY

We first observe the following topological fact which essentially says that the hypersurfaces Γ_0^ϵ from Theorem 4.6 are star-shaped relative to $\mathbf{0}$.

Proposition 5.1. *Let $\Sigma \subset \mathbb{R}^{n+1}$ be a connected hypersurface which is smoothly asymptotic to a regular cone $\mathcal{C}(\Sigma)$. If $\mathbf{x} \cdot \mathbf{n}_\Sigma > 0$ and $\Sigma \cap \mathcal{C}(\Sigma) = \emptyset$, then the map $\Pi : \Sigma \rightarrow \mathbb{S}^n$ given by*

$$(5.1) \quad \Pi(p) = \frac{\mathbf{x}(p)}{|\mathbf{x}(p)|}$$

is a diffeomorphism onto its image. Furthermore, the image is a connected component of $\mathbb{S}^n \setminus \mathcal{L}(\Sigma)$, where $\mathcal{L}(\Sigma)$ is the link of $\mathcal{C}(\Sigma)$.

Proof. Let $\Omega = \Pi(\Sigma)$ and observe that, as $\Pi(\mathcal{C}(\Sigma)) = \mathcal{L}(\Sigma)$ and $\Sigma \cap \mathcal{C}(\Sigma) = \emptyset$, $\Omega \cap \mathcal{L}(\Sigma) = \emptyset$. It follows that Π is a proper map. Indeed, suppose that $\mathcal{K} \subset \Omega$ is compact. As $\mathcal{K} \cap \mathcal{L}(\Sigma) = \emptyset$ and Σ is smoothly asymptotic to $\mathcal{C}(\Sigma)$, there is an $R = R(\mathcal{K}) > 0$ so that $\Pi^{-1}(\mathcal{K}) \subset B_R$. As Σ is a hypersurface, and so proper, $\Sigma \cap \bar{B}_R$ is compact and so $\Pi^{-1}(\mathcal{K}) \cap B_R$ is also compact.


Next notice that, as $\mathbf{x} \cdot \mathbf{n}_\Sigma > 0$ and

$$(5.2) \quad d\Pi_p(\mathbf{v}) = \frac{\mathbf{v}}{|\mathbf{x}(p)|} - \frac{\mathbf{x}(p)(\mathbf{x}(p) \cdot \mathbf{v})}{|\mathbf{x}(p)|^3} \quad \text{for } \mathbf{v} \in T_p\Sigma,$$

$d\Pi_p$ is bijective for all $p \in \Sigma$. Hence, Π is a local diffeomorphism and so Ω is an open subset of $\mathbb{S}^n$. As Ω is the image of a connected set, Ω itself must be connected and hence, by a standard topological result, Π is a finite covering map.

We now show that Π is a diffeomorphism. To see this, fix a $q \in \Omega$ and suppose that $p, p' \in \Pi^{-1}(q)$ are two consecutive points in the pre-image of q – here we order by distance from $\mathbf{0}$. For $\delta > 0$ sufficiently small, let $\tilde{p} = \mathbf{x}(p) + \delta\mathbf{x}(q)$ and likewise define $\tilde{p}'$. Invoking that $\mathbf{x} \cdot \mathbf{n}_\Sigma > 0$, the straight line segment $\tilde{p}\tilde{p}'$ connecting $\tilde{p}$ to $\tilde{p}'$ intersects transversally with Σ exactly once. Now let Π_Σ be the nearest point projection onto Σ and let $\gamma : [0, 1] \rightarrow \Sigma$ be a simple, smooth parametrized curve connecting $\Pi_\Sigma(\tilde{p})$ to $\Pi_\Sigma(\tilde{p}')$. Consider the curve $\tilde{\gamma} : [0, 1] \rightarrow \mathbb{R}^{n+1}$ defined by

$$(5.3) \quad \tilde{\gamma}(t) = \mathbf{x}(\gamma(t)) + \delta\mathbf{n}_\Sigma(\gamma(t)).$$


 FIGURE 1. The line segment $\overline{p\tilde{p}'}$ and curve $\tilde{\gamma}$.

Clearly, $\overline{p\tilde{\gamma}(0)} \cup \tilde{\gamma} \cup \overline{\tilde{\gamma}(1)p'}$ is homotopic to $\overline{p\tilde{p}'}$ in $\mathbb{R}^{n+1}$, but does not intersect Σ . This is a contradiction and completes the proof that Π is a diffeomorphism of Σ onto Ω .

Finally, we show that Ω must be a connected component of $\mathbb{S}^n \setminus \mathcal{L}(\Sigma)$. To see this, let $\hat{\Omega}$ be the connected component of $\mathbb{R}^{n+1} \setminus \mathcal{C}(\Sigma)$ which contains Σ – such a component exists as Σ is connected and disjoint from $\mathcal{C}(\Sigma)$. Clearly, $\Omega \subset \hat{\Omega} \cap \mathbb{S}^n$ and $\hat{\Omega} \cap \mathbb{S}^n$ is a connected component of $\mathbb{S}^n \setminus \mathcal{L}(\Sigma)$. Pick a point $q \in \partial\Omega$ and a sequence $q_i \in \Omega$ with $q_i \rightarrow q \notin \Omega$. Set $p_i = \Pi^{-1}(q_i) \in \Sigma$. As Σ is proper, $|\mathbf{x}(p_i)| \rightarrow \infty$ and so, as Σ is smoothly asymptotic to $\mathcal{C}(\Sigma)$, $q \in \mathcal{L}(\Sigma)$. Hence, $\partial\Omega \subset \mathcal{L}(\Sigma)$ and our claim is verified. $\square$

We may now prove Theorem 1.1.

Proof of Theorem 1.1. First observe that the theorem is trivially true for hyperplanes. We next claim that Σ must be connected. Indeed, otherwise one could use two of the connected components as barriers to find a stable self-shrinker which is impossible in view of [1, Lemma 3.3]. As such, we may assume that $\Sigma \in \mathcal{ACS}_n^*[\lambda_{n-1}]$.

Hence, applying Theorem 4.6, we associate to Σ two smooth mean curvature flows $\{\Gamma_t^{\pm\epsilon}\}_{t \in [-1, 0]}$ with $\Gamma_{-1}^{\pm\epsilon} = \Gamma^{\pm\epsilon}$ for some $\epsilon > 0$ small, where $\Gamma^{\pm\epsilon}$ are the hypersurfaces as in Proposition 4.2. Moreover, each $\Gamma^\pm = \Gamma_0^{\pm\epsilon}$ is a connected star-shaped hypersurface which is smoothly asymptotic to, but disjoint from, $\mathcal{C}(\Sigma)$. Notice that $\mathbb{R}^{n+1} \setminus \Sigma$ consists of exactly two connected components (as Σ is connected) one containing $\Gamma^{-\epsilon}$ and the other Γ^ϵ . Hence, $\mathbb{R}^{n+1} \setminus \mathcal{C}(\Sigma)$ also consists of exactly two connected components $\hat{\Omega}^\pm$ each

containing $\Gamma^\pm$ and so, by Proposition 5.1, each $\Gamma^\pm$ is diffeomorphic to $\Omega^\pm = \hat{\Omega}^\pm \cap \mathbb{S}^n$. This proves all but the last claim.

The final claim, that $\mathcal{L}(\Sigma)$ is connected, follows by applying the Mayer-Vietoris long exact sequence for reduced homology. Indeed, slightly enlarge $\Omega^\pm$ so that both are still connected and $\Omega^+ \cap \Omega^-$ is a regular tubular neighborhood of $\mathcal{L}(\Sigma)$ in $\mathbb{S}^n$. One then has

$$(5.4) \quad \tilde{H}_1(\mathbb{S}^n) \rightarrow \tilde{H}_0(\mathcal{L}(\Sigma)) \rightarrow \tilde{H}_0(\Omega^-) \oplus \tilde{H}_0(\Omega^+)$$

as part of the long exact sequence. As $n \geq 2$ and both Ω^- and Ω^+ are connected,

$$(5.5) \quad \{0\} = \tilde{H}_1(\mathbb{S}^n) = \tilde{H}_0(\Omega^-) = \tilde{H}_0(\Omega^+).$$

Hence, $\tilde{H}_0(\mathcal{L}(\Sigma)) = \{0\}$ and so $\mathcal{L}(\Sigma)$ is connected. $\square$

We next prove Corollary 1.2.

Proof of Corollary 1.2. First, notice that all possible singularities at the first singular time of mean curvature flow of closed surfaces have smooth supports – see [17, Theorem 2]. Hence, one can strengthen [9, Theorem 0.8] of Colding-Ilmanen-Minicozzi-White to conclude that any closed self-shrinker with entropy (equivalently, Gaussian surface area) less than or equal to λ_1 is diffeomorphic to a sphere. Furthermore, such a self-shrinker must be $\mathbb{S}_*^2$ by a recent result of Brendle [4] on the uniqueness of closed genus zero self-shrinkers (recall, in our definition self-shrinkers are automatically embedded).

Next, as observed in [2, Proposition 4.7] and in the proof of [2, Theorem 1.2] when $n = 2$, any non-compact self-shrinker in $\mathbb{R}^3$ with $\lambda[\Sigma] \leq \lambda_1$ is either a rotation of $\mathbb{S}_*^1 \times \mathbb{R}$ or asymptotically conical. By Theorem 1.1, in the latter case, Σ is diffeomorphic to an open disk. Invoking [4, Theorem 2] of Brendle, it follows that Σ must be flat. Thus, the only self-shrinkers with entropy less than or equal to λ_1 are $\mathbb{S}_*^2$ and rotations of $\mathbb{R}^2 \times \{0\}$ and $\mathbb{S}_*^1 \times \mathbb{R}$.

To prove the gap, we argue by contradiction. Namely, suppose that there was no such δ_0 . Then there would exist a sequence Σ_i of self-shrinkers with $\lambda[\Sigma_i] \in (\lambda_1, 2)$ and $\lambda[\Sigma_i] \rightarrow \lambda_1$. Up to passing to a subsequence, we may assume that Σ_i converges to a multiplicity one F -stationary varifold Σ with $\lambda[\Sigma] \leq \lambda_1$. As each Σ_i is two-sided, the tangent cones of Σ cannot be quasi-planes. Also, there is no minimal cone with isolated singularities in $\mathbb{R}^3$. Thus, by Allard's regularity theorem, Σ_i converges in $C_{loc}^\infty(\mathbb{R}^3)$ to Σ and $\lambda[\Sigma] = \lim_{i \rightarrow \infty} \lambda[\Sigma_i] = \lambda_1$. By the rigidity of self-shrinkers of small entropy established in the previous paragraph, this implies that (up to a rotation) $\Sigma = \mathbb{S}_*^1 \times \mathbb{R}$. Hence, in view of [8, Theorem 0.1], for all i sufficiently large, Σ_i are rotations of $\mathbb{S}_*^1 \times \mathbb{R}$ and, in particular, $\lambda[\Sigma_i] = \lambda_1$, giving the desired contradiction. $\square$

Finally we show Corollary 1.3.

Proof of Corollary 1.3. Let $\{\Sigma_t\}_{t \in [-1, T]}$ be the maximal smooth mean curvature flow with initial surface $\Sigma_{-1} = \Sigma$. As Σ is closed, a comparison with a large shrinking sphere implies that $T < \infty$. Let $X_0 = (\mathbf{x}_0, T)$ be a singular point of the flow. It follows from Corollary 1.2 and the monotonicity of entropy that $\lambda[\Sigma] \geq \lambda_2$ with equality if and only if, modulo translations and scalings, Σ is equal to $\mathbb{S}^2$. If the tangent flow at X_0 is a self-shrinking sphere, then, by Brakke's local regularity theorem, Σ_t must be very close to the sphere for all t near T . In particular, Σ_t is diffeomorphic to $\mathbb{S}^2$ and thus, as the flow is smooth, so is Σ . Hence, if Σ has positive genus, the tangent flow cannot be a shrinking sphere and so $\lambda[\Sigma] > \lambda_1$ by the monotonicity formula and Corollary 1.2. $\square$

APPENDIX A.

We use a weighted version of Huisken's monotonicity formula to establish a parabolic maximum principle for smooth non-compact mean curvature flows with boundaries. This is a slight generalization of [12, Corollary 1.1].

Theorem A.1. *Let $\mathcal{S} = \{\Sigma_t\}_{t \in [-1, T]}$ be a smooth mean curvature flow in $\mathbb{R}^{n+1}$ with finite entropy. For a fixed $R \geq 0$ and $X_0 = (\mathbf{0}, -1)$, suppose that u is a C^2 function on $\mathcal{S} \setminus C_{R, T+1}(X_0)$ which satisfies:*

(1) *on $\mathring{\mathcal{S}} = \mathcal{S} \setminus \overline{C_{R, T+1}(X_0)}$,*

$$(A.1) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) u \geq \mathbf{a} \cdot \nabla_{\Sigma_t} u + bu$$

with $\sup_{\mathcal{S}} |\mathbf{a}|^2 + |b| = M_0 < \infty$;

(2) *$u > c_0 \geq 0$ on $\partial_P \mathring{\mathcal{S}} = (\Sigma_{-1} \setminus B_R) \cup (\mathcal{S} \cap \partial C_{R, T+1}(X_0))$ for some constant c_0 ;*

(3) *for all $t \in [-1, T)$,*

$$(A.2) \quad \int_{\Sigma_t \setminus B_R} \left(|u|^2 + \left| \frac{du}{dt} \right|^2 + |\nabla_{\Sigma_t} u|^2 + |\nabla_{\Sigma_t}^2 u|^2 \right) \Phi_{(\mathbf{0}, T)} d\mathcal{H}^n < \infty,$$

where

$$(A.3) \quad \Phi_{(\mathbf{0}, T)}(\mathbf{x}, t) = (T - t)^{-\frac{n}{2}} \Phi \left(\frac{\mathbf{x}}{\sqrt{T - t}} \right) = (4\pi(T - t))^{-\frac{n}{2}} e^{-\frac{|\mathbf{x}|^2}{4(T - t)}}.$$

Then for all $t \in [-1, T)$,

$$(A.4) \quad \inf_{\Sigma_t \setminus B_R} u \geq c_0 e^{M_1(t+1)} \geq 0,$$

where $M_1 = \min \{ \inf_{\mathcal{S}} b, 0 \} \leq 0$.

Proof. We will first show that $u \geq 0$. Let $u_0(p, t) = \max \{-u(p, t), 0\} \geq 0$ on $\mathring{\mathcal{S}}$ and we extend u_0 by zero to the whole $\mathcal{S}$. As u is continuous and, for every $0 < T' < T$, $\mathcal{S} \cap \partial_P C_{R, T'+1}(X_0)$ is compact, Item (2) implies that $-u$ is strictly negative in a neighborhood of $\mathcal{S} \cap \partial_P C_{R, T'+1}(X_0)$. Hence, u_0^2 is $C^{1,1}$ and satisfies, in the weak sense, that

$$(A.5) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) u_0^2 \leq -2|\nabla_{\Sigma_t} u_0|^2 + 2u_0 \mathbf{a} \cdot \nabla_{\Sigma_t} u_0 + 2bu_0^2 \leq 3M_0 u_0^2.$$

Here we used Item (1) and Young's inequality.

Invoking (A.5) and Item (3), it follows from the weighted monotonicity formula [11, Theorem 4.13] that

$$(A.6) \quad \frac{d}{dt} \int_{\Sigma_t} u_0^2 \Phi_{(\mathbf{0}, T)} d\mathcal{H}^n \leq 3M_0 \int_{\Sigma_t} u_0^2 \Phi_{(\mathbf{0}, T)} d\mathcal{H}^n.$$

This together with Item (2) implies that for all $t \in [-1, T)$,

$$(A.7) \quad \int_{\Sigma_t} u_0^2 \Phi_{(\mathbf{0}, T)} d\mathcal{H}^n = 0.$$

That is, $u_0 \equiv 0$ on $\mathcal{S}$ and so $u \geq 0$ as claimed.

Set $\tilde{u} = e^{-M_1(t+1)} u$. Using $\tilde{u} \geq 0$ and $b - M_1 \geq 0$, it follows that

$$(A.8) \quad \left(\frac{d}{dt} - \Delta_{\Sigma_t} \right) \tilde{u} \geq \mathbf{a} \cdot \nabla_{\Sigma_t} \tilde{u} + (b - M_1) \tilde{u} \geq \mathbf{a} \cdot \nabla_{\Sigma_t} \tilde{u}.$$

As $M_1 \leq 0$, $\tilde{u} > c_0$ on $\partial_P \mathring{S}$. Applying the same argument as above to $\tilde{u} - c_0$, we conclude that $\tilde{u} \geq c_0$ which completes the proof. $\square$

APPENDIX B.

Throughout this appendix, let Σ be a self-shrinker in $\mathbb{R}^{n+1}$ and let $H_w^1(\Sigma)$ and $L_w^2(\Sigma)$ be as in the proof of Proposition 4.1. The goal is to prove the natural embedding

$$(B.1) \quad \iota : H_w^1(\Sigma) \hookrightarrow L_w^2(\Sigma)$$

is compact.

First, we prove a lemma that yields a uniform decay rate for the weighted L^2 integration over exterior regions.

Lemma B.1. *For all $\phi \in C_c^\infty(\Sigma)$,*

$$(B.2) \quad \int_{\Sigma} \phi^2 |\mathbf{x}|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n \leq 16 \int_{\Sigma} |\nabla_{\Sigma} \phi|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n + 4n \int_{\Sigma} \phi^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n$$

Proof. Recall that the divergence theorem implies that

$$(B.3) \quad \int_{\Sigma} \operatorname{div}_{\Sigma} \mathbf{V} d\mathcal{H}^n = - \int_{\Sigma} \mathbf{H}_{\Sigma} \cdot \mathbf{V} d\mathcal{H}^n$$

for all vector fields $\mathbf{V}$ of compact support. Setting $\mathbf{V} = \mathbf{x} \phi^2 e^{-|\mathbf{x}|^2/4}$ yields the lemma.

Indeed, we compute

$$(B.4) \quad \operatorname{div}_{\Sigma} \mathbf{V} = n \phi^2 e^{-\frac{|\mathbf{x}|^2}{4}} + 2(\mathbf{x} \cdot \nabla_{\Sigma} \phi) \phi e^{-\frac{|\mathbf{x}|^2}{4}} - \frac{1}{2} \phi^2 |\mathbf{x}^{\top}|^2 e^{-\frac{|\mathbf{x}|^2}{4}},$$

and, using the self-shrinker equation and Young's inequality,

$$(B.5) \quad \begin{aligned} \frac{1}{2} \int_{\Sigma} \phi^2 |\mathbf{x}|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n &= n \int_{\Sigma} \phi^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n + 2 \int_{\Sigma} (\mathbf{x} \cdot \nabla_{\Sigma} \phi) \phi e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n \\ &\leq n \int_{\Sigma} \phi^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n + 4 \int_{\Sigma} |\nabla_{\Sigma} \phi|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n \\ &\quad + \frac{1}{4} \int_{\Sigma} \phi^2 |\mathbf{x}|^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n. \end{aligned}$$

The desired estimate follows from a rearrangement of the above inequality. $\square$

Now we are ready to prove that

Proposition B.2. *The natural embedding $\iota : H_w^1(\Sigma) \hookrightarrow L_w^2(\Sigma)$ is compact.*

Proof. As $C_c^\infty(\Sigma)$ is dense in $H_w^1(\Sigma)$, it suffices to prove the proposition for ι restricted to $C_c^\infty(\Sigma) \cap H_w^1(\Sigma)$. Namely, let ϕ_i be a sequence of elements of $C_c^\infty(\Sigma) \cap H_w^1(\Sigma)$ with $(\phi_i, \phi_i)_1 \leq C$ for some constant $C > 0$. We need to find a subsequence ϕ_{i_j} converging strongly in $L_w^2(\Sigma)$.

For $R > 1$ we introduce the cut-off function $\psi_R : \mathbb{R}^{n+1} \rightarrow [0, 1]$ so that $\psi_R = 1$ in B_R , $\psi_R = 0$ outside B_{2R} , and $|D\psi_R| \leq 2/R$. Let $\phi_i^R = \phi_i \psi_R$. Clearly $\operatorname{spt}(\phi_i^R) \subset \Sigma \cap B_{2R}$ and $(\phi_i^R, \phi_i^R)_1 \leq 8C$ for all i . By the Rellich-Kondrachov compactness theorem, there is a subsequence of ϕ_i^R converging strongly in $L_w^2(\Sigma)$ to ϕ^R supported in $\Sigma \cap B_{2R}$. Now take an increasing sequence $R_k \rightarrow \infty$ and by a diagonalization argument we find a subsequence $i_j \rightarrow \infty$ and $\phi \in L_w^2(\Sigma)$ so that $\phi_{i_j} \rightarrow \phi$ strongly in $L_w^2(\Sigma \cap B_{R_k})$ for each k .

Observe that $\|\phi^{R_k} - \phi\|_0 \rightarrow 0$ as $k \rightarrow \infty$. And by Lemma B.1, $\|\phi_{i_j} - \phi_{i_j}^{R_k}\|_0 \leq C' R_k^{-2}$ for some C' depending only on n and C . Thus

$$(B.6) \quad \|\phi_{i_j} - \phi\|_0 \leq \|\phi_{i_j} - \phi_{i_j}^{R_k}\|_0 + \|\phi_{i_j}^{R_k} - \phi^{R_k}\|_0 + \|\phi^{R_k} - \phi\|_0 \rightarrow 0,$$

i.e., $\phi_{i_j} \rightarrow \phi$ strongly in $L_w^2(\Sigma)$. $\square$

APPENDIX C.

Let $\Sigma \in \mathcal{ACS}_n^*$ and let f be the lowest eigenfunction of L_Σ as in Proposition 4.1. Throughout $\epsilon_1 > 0$ is assumed to be sufficiently small. We consider a one-parameter family of hypersurfaces

$$(C.1) \quad \Gamma^s = \{\mathbf{y} \in \mathbb{R}^{n+1} : \mathbf{y} = \mathbf{x}(p) + sf(p)\mathbf{n}_\Sigma(p), p \in \Sigma\}$$

for $|s| \leq \epsilon_1$ with $\Gamma^0 = \Sigma$. Observe that, as Γ^s is asymptotically conical by Item (3) of Proposition 4.2, for each $|s| \leq \epsilon_1$, this family of hypersurfaces is a normal variation of Γ^s with vector field $f_{\Gamma^s} \mathbf{n}_{\Gamma^s}$, where $f_{\Gamma^s} = f \mathbf{n}_\Sigma \cdot \mathbf{n}_{\Gamma^s}$. Define a function

$$(C.2) \quad G : \mathbb{R}^{n+1} \times \mathbb{R}^+ \times [-\epsilon_1, \epsilon_1] \rightarrow \mathbb{R}^+, \quad G(\mathbf{x}_0, t_0, s) = (4\pi t_0)^{-\frac{n}{2}} \int_{\Gamma^s} e^{-\frac{|\mathbf{x} - \mathbf{x}_0|^2}{4t_0}} d\mathcal{H}^n.$$

The goal of this appendix is to show that there is an $\epsilon_2 \in (0, \epsilon_1)$ sufficiently small so that if $s \neq 0$ and $|s| \leq \epsilon_2$, then

$$(C.3) \quad \lambda[\Gamma^s] \equiv \sup_{\mathbf{x}_0, t_0} G(\mathbf{x}_0, t_0, s) < G(\mathbf{0}, 1, 0) = \lambda[\Sigma].$$

In order to establish (C.3), we follow closely the proof of [10, Theorem 0.15] and thus it suffices to show:

Proposition C.1. *The following properties hold:*

- (1) G has a strict local maximum at $(\mathbf{0}, 1, 0)$;
- (2) The restriction of G to Σ , i.e., $G(\mathbf{x}_0, t_0, 0)$, has a strict global maximum at $(\mathbf{0}, 1)$;
- (3) $|\partial_s G|$ is uniformly bounded on compact sets;
- (4) $G(\mathbf{x}_0, t_0, s)$ is strictly less than $G(\mathbf{0}, 1, 0)$ whenever $|\mathbf{x}_0|$ is sufficiently large;
- (5) $G(\mathbf{x}_0, t_0, s)$ is strictly less than $G(\mathbf{0}, 1, 0)$ whenever $|\log t_0|$ is sufficiently large.

Proof. Observe that, by [10, Proposition 3.6], the gradient of G vanishes at $(\mathbf{0}, 1, 0)$. Hence, Item (1) will follow if we show that the Hessian of G at $(\mathbf{0}, 1, 0)$ is negative definite. That is, if, for any $a, b \in \mathbb{R}$ and $\mathbf{y} \in \mathbb{R}^{n+1}$ so that $(a, b, \mathbf{y}) \neq (0, 0, \mathbf{0})$,

$$(C.4) \quad \partial_{ss}|_{s=0} G(s\mathbf{y}, 1 + as, bs) < 0.$$

To see this, we first use [10, Theorem 4.14] to conclude that, at $s = 0$,

$$(C.5) \quad \begin{aligned} \partial_{ss} G(s\mathbf{y}, 1 + as, s) &= (4\pi)^{-\frac{n}{2}} \int_{\Sigma} \left(-f L_\Sigma f - a^2 H_\Sigma^2 - \frac{(\mathbf{y} \cdot \mathbf{n}_\Sigma)^2}{2} \right) e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n \\ &\quad + (4\pi)^{-\frac{n}{2}} \int_{\Sigma} (2af H_\Sigma + f \mathbf{y} \cdot \mathbf{n}_\Sigma) e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n. \end{aligned}$$

Note that f , H_Σ , and $\mathbf{y} \cdot \mathbf{n}_\Sigma$ are eigenfunctions corresponding to different eigenvalues of L_Σ . As Σ is asymptotically conical and f satisfies (4.2) and (4.3), it follows from [10,

Corollary 3.10] that f is orthogonal to both H_Σ and $\mathbf{y} \cdot \mathbf{n}_\Sigma$ in $L_w^2(\Sigma)$. This implies that the last integration in the right hand side of (C.5) is zero and so, at $s = 0$,

$$(C.6) \quad \begin{aligned} \partial_{ss} G(s\mathbf{y}, 1 + as, s) &= (4\pi)^{-\frac{n}{2}} \int_\Sigma \left(-f L_\Sigma f - a^2 H_\Sigma^2 - \frac{(\mathbf{y} \cdot \mathbf{n}_\Sigma)^2}{2} \right) e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n \\ &\leq \mu(4\pi)^{-\frac{n}{2}} \int_\Sigma f^2 e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n < 0. \end{aligned}$$

Hence, for any $b \in \mathbb{R} \setminus \{0\}$, the second derivative $\partial_{ss}|_{s=0} G(s\mathbf{y}, 1 + sa, bs) < 0$. Finally, appealing to [10, Theorem 4.14] again,

$$(C.7) \quad \partial_{ss}|_{s=0} G(s\mathbf{y}, 1 + sa, 0) = (4\pi)^{-\frac{n}{2}} \int_\Sigma \left(-a^2 H_\Sigma^2 - \frac{(\mathbf{y} \cdot \mathbf{n}_\Sigma)^2}{2} \right) e^{-\frac{|\mathbf{x}|^2}{4}} d\mathcal{H}^n < 0,$$

where the inequality follows from the fact that $\Sigma \in \mathcal{ACS}_n^*$ and so Σ can neither split off a line isometrically (and so is non-flat) nor can H_Σ vanish identically. Hence, we conclude that the Hessian of G at $(\mathbf{0}, 1, 0)$ is negative definite as claimed. In particular, there is an $\epsilon_3 \in (0, \epsilon_1)$ so that

$$(C.8) \quad G(\mathbf{x}_0, t_0, s) < G(\mathbf{0}, 1, 0) \quad \text{if } 0 < |\mathbf{x}_0|^2 + (\log t_0)^2 + s^2 < \epsilon_3^2.$$

Next, [10, Lemma 7.10] implies that $\lambda[\Sigma] = G(\mathbf{0}, 1, 0)$, and there is an $\alpha_0 > 0$ so that

$$(C.9) \quad G(\mathbf{x}_0, t_0, 0) < G(\mathbf{0}, 1, 0) - \alpha_0 \quad \text{if } |\mathbf{x}_0|^2 + (\log t_0)^2 > \frac{\epsilon_3^2}{4},$$

which gives Item (2).

As $\lambda[\Sigma] < \infty$, Σ has bounded area ratios. Since Σ has uniformly bounded second fundamental form, combining with (4.2) and (4.3), it follows from [25, (2.34)] that

$$(C.10) \quad \sup_{|s| \leq \epsilon_1} \sup_{\mathbf{x} \in \mathbb{R}^{n+1}, r > 1} r^{-n} \mathcal{H}^n(\Gamma^s \cap B_r(\mathbf{x})) < C(\Sigma, C_0, C_1).$$

Furthermore, as Σ is asymptotically conical, by (4.2) and (4.3), there is a C_H depending only on C_0, C_1, C_2 , and Σ so that

$$(C.11) \quad \sup_{|s| \leq \epsilon_1} \sup_{\Gamma^s} H_{\Gamma^s}^2 \leq C_H.$$

Then, using the first variation formula [10, Lemma 3.1], we argue as in [10, (7.5)] to get that

$$(C.12) \quad \partial_{t_0} G(\mathbf{x}_0, t_0, s) \geq -(4\pi t_0)^{-\frac{n}{2}} \int_{\Gamma^s} \frac{H_{\Gamma^s}^2}{4} e^{-\frac{|\mathbf{x} - \mathbf{x}_0|^2}{4t_0}} d\mathcal{H}^n \geq -\frac{C_H}{4} G(\mathbf{x}_0, t_0, s).$$

Thus, for $0 < t_0 \leq \bar{t}$,

$$(C.13) \quad G(\mathbf{x}_0, t_0, s) \leq e^{\frac{C_H \bar{t}}{4}} G(\mathbf{x}_0, \bar{t}, s).$$

In particular, choosing $\bar{t} = 1$, this together with (C.10) implies that

$$(C.14) \quad \sup_{|s| \leq \epsilon_1} \sup_{\mathbf{x} \in \mathbb{R}^{n+1}, r > 0} r^{-n} \mathcal{H}^n(\Gamma^s \cap B_r(\mathbf{x})) < C_M,$$

where C_M depends only on n, C , and C_H .

An easy consequence of (C.14) is that given $\alpha_1 > 0$ small, there is an $R > 0$ so that

$$(C.15) \quad \sup_{|s| \leq \epsilon_1} \sup_{\mathbf{x}_0, t_0} (4\pi t_0)^{-\frac{n}{2}} \int_{\Gamma^s \setminus B_{R\sqrt{t_0}}(\mathbf{x}_0)} \frac{|\mathbf{x} - \mathbf{x}_0|}{\sqrt{t_0}} e^{-\frac{|\mathbf{x} - \mathbf{x}_0|^2}{4t_0}} d\mathcal{H}^n < \frac{\alpha_1}{4}.$$

Invoking the first variation formula again, we get that

$$(C.16) \quad \partial_s G(\mathbf{x}_0, t_0, s) = (4\pi t_0)^{-\frac{n}{2}} \int_{\Gamma^s} f_{\Gamma^s} \left(H_{\Gamma^s} - \frac{(\mathbf{x} - \mathbf{x}_0) \cdot \mathbf{n}_{\Gamma^s}}{2t_0} \right) e^{-\frac{|\mathbf{x} - \mathbf{x}_0|^2}{4t_0}} d\mathcal{H}^n.$$

Let $\mathcal{K}$ be any compact set in $\mathbb{R}^{n+1} \times \mathbb{R}^+ \times [-\epsilon_1, \epsilon_1]$. There is an $\bar{R}$ so that $B_{\bar{R}}$ contains $B_{R\sqrt{t_0}}(\mathbf{x}_0)$ for every $(\mathbf{x}_0, t_0, s) \in \mathcal{K}$. Thus, with (4.2) and (C.11), it follows from (C.15) that the integration in (C.16) restricted on $\mathbb{R}^{n+1} \setminus B_{\bar{R}}$ will be uniformly bounded on $\mathcal{K}$. Clearly, the integration in (C.16) restricted on $B_{2\bar{R}}$ is continuous in all three variables $\mathbf{x}_0$, t_0 , and s . Hence we conclude that $\partial_s G$ is uniformly bounded on $\mathcal{K}$, proving Item (3).

To see Item (4), observe that, by Item (3) of Proposition 4.2, for any given $\alpha_1 > 0$ small, there is an $R' > 1$ so that if $|\mathbf{x}_0| > R'$, $0 < t_0 \leq 1$, and $|s| \leq \epsilon_1$, then either $(\Gamma^s - \mathbf{x}_0)/\sqrt{t_0}$ is contained outside $B_{R'}$, or $(\Gamma^s - \mathbf{x}_0)/\sqrt{t_0}$ in $B_{2R'}$ is a connected small C^1 graph over some hyperplane. This, together with (C.15), implies that $G(\mathbf{x}_0, t_0, s) < 1 + \alpha_1/2$.

Next we study the region where $t_0 > 1$ and $|\mathbf{x}_0|$ sufficiently large. Set $\rho = 1/\sqrt{t_0} < 1$ and $\mathbf{y}_0 = \mathbf{x}_0/\sqrt{t_0}$. By (C.14), given $\alpha_1 > 0$ small, there is an $r_1 > 0$ small so that for all $\mathbf{y}_0 \in \mathbb{R}^{n+1}$, $\rho < 1$, and $|s| \leq \epsilon_1$,

$$(C.17) \quad (4\pi)^{-\frac{n}{2}} \int_{\rho\Gamma^s \cap B_{r_1}} e^{-\frac{|\mathbf{x} - \mathbf{y}_0|^2}{4}} d\mathcal{H}^n < \frac{\alpha_1}{4}.$$

Observe that $\rho\Gamma^s$ is the normal graph of the function $s\rho f(\rho^{-1}\cdot)$ on $\rho\Sigma$. As $\rho\Sigma \rightarrow \mathcal{C}(\Sigma)$ in $C_{loc}^\infty(\mathbb{R}^{n+1} \setminus \{\mathbf{0}\})$ as $\rho \rightarrow 0$ and the link $\mathcal{L}(\Sigma)$ is smooth embedded, it follows from (4.2) and (4.3) that there is a $C_A(r_1)$ so that

$$(C.18) \quad \sup_{|s| \leq \epsilon_1, \rho < 1} \sup_{\rho\Gamma^s \setminus B_{r_1}} |A_{\rho\Gamma^s}| < C_A(r_1).$$

Furthermore, together with (C.14), (C.15) and (C.17), we get that, if $\mathbf{y}_0 \in \mathbb{R}^{n+1}$, $\rho < 1$, and $|s| \leq \epsilon_1$, then

$$(C.19) \quad \begin{aligned} F[\rho\Gamma^s - \mathbf{y}_0] &\leq F[\rho\Sigma - \mathbf{y}_0] + \frac{\alpha_1}{2} + (4\pi)^{-\frac{n}{2}} \int_{\rho\Gamma^s \cap (B_R(\mathbf{y}_0) \setminus B_{r_1})} e^{-\frac{|\mathbf{x} - \mathbf{y}_0|^2}{4}} d\mathcal{H}^n \\ &\quad - (4\pi)^{-\frac{n}{2}} \int_{\rho\Sigma \cap (B_R(\mathbf{y}_0) \setminus B_{r_1})} e^{-\frac{|\mathbf{x} - \mathbf{y}_0|^2}{4}} d\mathcal{H}^n \\ &\leq F[\rho\Sigma - \mathbf{y}_0] + \frac{\alpha_1}{2} + C_E, \end{aligned}$$

where C_E depends on $C_0, C_1, \epsilon_1, r_1, C_M$, and C_A with $C_E \rightarrow 0$ as $\epsilon_1 \rightarrow 0$.

Now let us choose $\alpha_1 = \min\{\alpha_0/4, 2(\lambda[\Sigma] - 1)/5\}$ and possibly shrink ϵ_1 so that $C_E < \alpha_0/4$. Hence, by (C.9), we conclude that for all $|\mathbf{x}_0| > \bar{R}' = \max\{R', \frac{\epsilon_3}{2}\}$

$$(C.20) \quad G(\mathbf{x}_0, t_0, s) < \lambda[\Sigma] - \frac{\alpha_0}{2}.$$

Finally, in view of (C.19), we only need to deal with the case that t_0 approaches to 0 and $|\mathbf{x}_0| \leq \bar{R}'$ so to complete the proof of Item (5). We choose $\bar{t}$ sufficiently small so that

$$(C.21) \quad e^{\frac{C_H \bar{t}}{4}} (\lambda[\Sigma] - \alpha_0) < \lambda[\Sigma] - \frac{\alpha_0}{2}.$$

Thus, further shrinking ϵ_1 if necessary, it follows from (C.13), (C.9) and Item (3) that for all $0 < t_0 < \bar{t}$, $|\mathbf{x}_0| \leq \bar{R}'$, and $|s| \leq \epsilon_1$, we have that $G(\mathbf{x}_0, t_0, s) < \lambda[\Sigma] - \alpha_0/4$. $\square$

Acknowledgements. We would like to thank Detang Zhou for many constructive comments, in particular for explaining to us an alternative proof of the half-space theorem for self-shrinkers and the Rellich-Kondrachov type theorem of weighted Sobolev spaces.

REFERENCES

- [1] J. Bernstein and L. Wang, *A remark on a uniqueness property of high multiplicity tangent flows in dimension three*, Int. Math. Res. Not. IMRN 2015, no. 15, 6286–6294.
- [2] J. Bernstein and L. Wang, *A sharp lower bound for the entropy of closed hypersurfaces up to dimension six*, preprint. Available at: <http://arxiv.org/abs/1406.2966>.
- [3] K. Brakke, *The motion of a surface by its mean curvature*, Mathematical Notes 20, Princeton University Press, Princeton, N.J., 1978.
- [4] S. Brendle, *Embedded self-similar shrinkers of genus 0*, Ann. of Math. (2) 183, (2016), no. 2, 715–728.
- [5] M. Cavalcante and J. Espinar, *Halfspace type theorems for self-shrinkers*, to appear in Bull. London Math. Soc.. Available at: <http://arxiv.org/abs/1412.3754>.
- [6] B. L. Chen and L. Yin, *Uniqueness and pseudolocality theorems of the mean curvature flow*, Comm. Anal. Geom. 15 (2007), no. 3, 435–490.
- [7] X. Cheng and D. Zhou, *Eigenvalues of the drifted Laplacian on complete metric measure spaces*, to appear in Commun. Contemp. Math.. Available at: <http://arxiv.org/abs/1305.4116>.
- [8] T.H. Colding, T. Ilmanen, and W.P. Minicozzi II, *Rigidity of generic singularities of mean curvature flow*, Publ. Math. Inst. Hautes Études Sci. 121 (2015), no. 1, 363–382.
- [9] T.H. Colding, T. Ilmanen, W.P. Minicozzi II, and B. White, *The round sphere minimizes entropy among closed self-shrinkers*, J. Differential Geom. 95 (2013), no. 1, 53–69.
- [10] T. H. Colding and W.P. Minicozzi II, *Generic mean curvature flow I: generic singularities*, Ann. of Math. (2) 175 (2012), no. 2, 755–833.
- [11] K. Ecker, *Regularity theory for mean curvature flow*, Progress in Nonlinear Differential Equations and their Applications, 57, Birkhäuser Boston, Inc., Boston, MA, 2004.
- [12] K. Ecker and G. Huisken, *Mean curvature evolution of entire graphs*, Ann. of Math. (2) 130 (1989), no. 3, 453–471.
- [13] K. Ecker and G. Huisken, *Interior estimates for hypersurfaces moving by mean curvature*, Invent. Math. 105 (1991), no. 3, 547–569.
- [14] D. Gilbarg and N. Trudinger, *Elliptic partial differential equations of second order*, Second edition, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 224, Springer-Verlag, Berlin, 1983.
- [15] G. Huisken, *Local and global behaviour of hypersurfaces moving by mean curvature*, Differential geometry: partial differential equations on manifolds (Los Angeles, CA, 1990), 175–191, Proc. Sympos. Pure Math., 54, Part 1, Amer. Math. Soc., Providence, RI, 1993.
- [16] G. Huisken, *Asymptotic behaviour for singularities of the mean curvature flow*, J. Differential Geom. 31 (1990), no. 1, 285–299.
- [17] T. Ilmanen, *Singularities of mean curvature flow of surfaces*, preprint. Available at: <http://www.math.ethz.ch/~ilmanen/papers/pub.html>.
- [18] T. Ilmanen, A. Neves, and F. Schulze, *On short time existence for the planar network flow*, preprint. Available at: <http://arxiv.org/abs/1407.4756>.
- [19] G.M. Lieberman, *Second order parabolic differential equations*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [20] H. Samelson, *Orientability of hypersurfaces in $\mathbb{R}^n$* , Proc. Amer. Math. Soc. 22 (1969), 301–302.
- [21] N. Sesum, *Rate of convergence of the mean curvature flow*, Comm. Pure Appl. Math. 61 (2008), no. 4, 464–485.
- [22] L. Simon, *Existence of surfaces minimizing the Willmore functional*, Comm. Anal. Geom. 1 (1993), no. 2, 281–326.
- [23] K. Smoczyk, *Starshaped hypersurfaces and the mean curvature flow*, Manuscripta Math. 95 (1998), no. 2, 225–236.
- [24] A. Stone, *A density function and the structure of singularities of the mean curvature flow*, Calc. Var. Partial Differential Equations 2 (1994), no. 4, 443–480.
- [25] L. Wang, *Uniqueness of self-similar shrinkers with asymptotically conical ends*, J. Amer. Math. Soc. 27 (2014), no. 3, 613–638.
- [26] B. White, *A local regularity theorem for mean curvature flow*, Ann. of Math. (2) 161 (2005), no. 3, 1487–1519.

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, 3400 N. CHARLES STREET, BALTIMORE, MD 21218

E-mail address: `bernstein@math.jhu.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, 480 LINCOLN DRIVE, MADISON, WI 53706

E-mail address: `luwang@math.wisc.edu`